Plant Disease Reporter

Agricultural Research Center

Agricultural Research Service

UNITED STATES DEPARTMENT OF AGRICULTURE

VOLUME 56 NUMBER 13

1972 Index
(Issued June 1973)

The Plant Disease Reporter is published monthly as a service to American agriculture. The articles emphasize new developments in plant pathology, including disease incidence and severity, serious outbreaks and epidemics, investigative techniques, and control chemicals. The rapid issuance of such information permits plant pathologists to recommend the judicious use of pesticides or other appropriate control measures against impending disease epidemics. Agricultural Research Center, in publishing this material, does not assume responsibility for the subject matter.

GUIDE TO MANUSCRIPT PREPARATION

GENERAL: Consult recent issues of PLANT DISEASE REPORTER. Follow the journal format. Manuscripts must be the original ribbon copy. Length of articles is limited to 12 pages typed double-spaced throughout, including tables, graphs, and photographs. Author's name should appear in top right-hand corner of each page. Submission implies nonsubmission elsewhere and (if accepted) no publication elsewhere in the same form without consent.

SUBMISSION OF MANUSCRIPTS: Submit all manuscripts to: Plant Disease Reporter, Applied Plant Pathology Laboratory, Agricultural Research Center, Beltsville, Maryland 20705.

TITLE AND ABSTRACTS: Titles should be brief and informative. Abstracts are requested for all except very short papers. Additional Key Words, not in the Title or Abstract, may be added at the end of the Abstract.

CAUSES OF DISEASES: Give Latin names and authorities for pathogens causing parasitic diseases. Do not give authorities for host plants. For viruses, use either an accepted common name or a Latin name. For nonparasitic diseases, state the causal factor if it is known.

LITERATURE CITED: Alphabetize and number for citation in text. All references should be cited in text, and agree with Literature Cited numbers. Place year of publication after author's name. Cite number or month for periodicals that are not paged consecutively throughout the volume. Please check your references carefully, because we cannot always do it for you.

PESTICIDES: At the first mention of a pesticide or other agricultural chemical, or complex compound, use the full chemical name followed by the accepted common name. Thereafter, use the common name. If there is no common name, use an acceptable designation. Do not use a trade or brand name unless there is no other. Trade names in USDA papers must be footnoted with the official USDA disclaimer.

ILLUSTRATIONS: Keep figures to a minimum and do not mount them. When necessary, mark the "top" of each print and write figure numbers lightly on the back in pencil. Line cuts may be photographs or originals, but should be finished and ready for reproduction. Please keep in mind that the Reporter page before reproduction measures 8 inches wide by 11 1/2 inches long, and is reduced 25% in the photographic process. Figures are also reduced 25%, so be sure that the ones you submit are not too small.

TABLES AND GRAPHS: Make the titles and headings of tables, and the legends for graphs, definite and self-explanatory. Designate footnotes for tables with superscript lower-case letters. Be sure that text discussion agrees with data in the tables. Lettering in graphs should be clear and large enough to be legible after reduction. Insofar as possible, data should be presented as graphs rather than tables.

NEGATIVES: After publication, the negatives of each paper will be mailed automatically to the author or his designated representative. These negatives may be used to make the multilith plates from which reprints can be run. If reprints are desired of articles other than your own, please request them directly from the authors. Material published in the PLANT DISEASE REPORTER is public property; however, permission to reproduce illustrations or data appearing in it should be obtained from the author of the paper, with an appropriate credit line to PDR as the original source of publication.

SUBJECT MATTER

Subject matter should emphasize new things in plant pathology, such as records of disease occurrence, serious outbreaks and epidemics, conditions affecting development of plant diseases, techniques of investigation including instrumentation, discoveries in control including new materials and their evaluation. Papers cannot be accepted for publication that report routine control experiments, reviews, bibliographies without annotation, routine surveys, mere summaries or lists of plant diseases, original descriptions of new species, reports of insect damage, and papers dealing primarily with mycology, genetics, or physiology.

RECIPIENTS OF THE PLANT DISEASE REPORTER

The Plant Disease Reporter will be sent free to cooperating scientists in the Department of Agriculture and its associated Agencies who are engaged in plant pathological work, such as Federal and State research and extension plant pathologists. Any person who is not eligible to receive this publication without charge may subscribe to it by writing to: Superintendent of Documents, U. S. Government Printing Office, Washington, D.C. 20402.

PLANT DISEASE REPORTER INDEX

Issued by

APPLIED PLANT PATHOLOGY LABORATORY
Plant Protection Institute

Agricultural Research Center
Volume 56

Beltsville, Maryland
Number 13

AUTHOR INDEX

AAPOLA, ALPO, 198
ABAWI, G. S., (434)
ABDOU, Y. A., (878)
ABOUL-EID, H. Z., 699
ADEGBOLA, M. O. K., 755
ADLERZ, W. C., 563
AFFELTRANGER, CHARLES E., 457
ALBREGTS, E. E., (23)
ALCONERO, RODRIGO, 320, 501
ALDWINCKLE, HERB S., 616
ALFIERI, S. A., Jr., 511
ALLEN, D. J., 678
ALTMAN, JACK, 1067
ARAGAKI, M., 73
ARNESON, P. A., (616)
ARNETT, JAMES D., 310
ARTHURTON, P. A. J., (817)
ARTMAN, J. D., 66, 958
ATKINS, JOHN G., 26
AULAKH, KIRPAL S., (1099)
AYERS, J. E., 836
AZAD, H. R., (206)
AZERI, TURHAN, 352

BACKMAN, P. A., (362)
BAGGETT, J. R., 131
BAIN, D. C., 210
BAMMI, R. K., 990
BANIHASHEMI, Z., 206
BAR-JOSEPH, M., (1035)
BARKSDALE, T. H., 321
BARNES, GEORGE L., 35, 980, 1022
BARRITT, B. H., (792)
BEARD, J. B., (32)
BECKER, R. F., (434)
BENDER, FREDERICK W., (523)
BENNETT, J. E., (1069)
BENNETT, O. L., 371
BERG, C. C., (75)
BERGER, R. D., (661)
BERGQUIST, R. R., 112, 329
BERRY, CHARLES D., 614
BERRY, J. W., Jr., (398)
BHARGAVA, S. N., 64
BIELENBERG, O., (286)
BIRCHFIELD, W., (58)
BLANCO, M. H., 889
BLAZQUEZ, C. H., 243
BLOUNT, V. L., (65), (237), (268)
BLOWERS, D. R., (39)
BODE, F. L., (572)
BOHN, G. W., (935)
BOUWKAMP, JOHN C., (783)
BRADFUTE, O. E., (652)
BRASHEARS, ALAN D., (1065)
BRATHWAITE, C. W. D., 618, 1077
BRAVERMAN, SAMUEL W., 618
BRENNAN, EILEEN, 85
BROADBENT, P., 80
BRÖNNIMANN, A., 188, (761)
BROWDER, L. E., 847
BROWN, A. R., (534)
BROWN, G. ELDON, 909
BROWN, GUY E., 59
BROWN, J. F., (597)
BRUEHL, G. W., 20, 594
BURCHILL, R. T., (770)
BURGIS, D. S., (504), (695)
BURNSIDE, K. R., (368), (515)
BUTTERFIELD, J. E., (88)
BYRD, W. P., (169)

CALAVAN, E. C., 976
CALLIHAN, R. H., (400)
CAMERON, J. W., 927
CAMPBELL, J. M., (708)
CARVER, R. B., (69), (157), (368), (515), (622)
CEPONIS, M. J., 88
CHAHAL, AJAIB S., 601, 1099
CHALKLEY, JAMES H., (147)
CHANG, HELEN Y., 1028
CHEO, P. C., 1010
CHOHAN, J. S., (115)
CHRISTENSEN, C. M., 173, 1097
CHRISTIE, W. D., (53)
CLONINGER, F. D., (11)
COLE, H., Jr., (6), (59), (164), (386)
COMPTON, W. A., (863)
COMSTOCK, J. C., 247
CONNOR, M. D., (405)
CONVERSE, R. H., 1007
COOK, MARION, 382
COOK, R., 261
COX, R. S., 656
CRANDALL, BOWEN S., 916, 917, 1047, 1049
CRANDALL, P. C., (467)
CRAWFORD, JOHNNY L., 328
CRILL, PAT, 504, 695, 817
CRITCHLEY, P., (869)
CUNFER, B., (20)

DAFT, GILBERT C., (480)
DAINELLO, F. J., (957)
DAINES, ROBERT H., 122
DAKWA, J. T., 71, 1011
DALE, J. L., 270, 355, 658
DALE, P. S., 850
DARLINGTON, LEE C., (788)
DAUBENY, HUGH A., 792
DAVIDSON, ROY M., Jr., 565
DAVIS, J. R., 400
DAVIS, TERRY C., (108), 192
DAVISON, ARLEN D., 50, (1007)
DAVISON, E. M., 290
DELBOS, R., (293)
DEMSKI, JAMES W., (146), 147
DENMARK, J. C., (511)
DERRICK, K. S., (69), (622)
DeWOLFE, T. A., (927)
DIAZ, FRANCISCO FISCHNALER, 437
DICKERSON, O. J., 957
DICKSON, R. C., (722)
DOCHINGER, LEON S., 523
DOHNER, L. D., (45)
DOLAN, DESMOND D., (618)
DOOLEY, H. L., 1069
DOUGLAS, DEXTER R., 308
DRIVER, CHAS. H., (407)
DUAFALA, T., (313)
DuCHARME, E. P., 85
DUNEZ, J., 293
DUNN, R. A., 577
DURBIN, R. D., (949)

EBBEN, MARION H., 869
EBRON, SONIA P., (960)
ECHANDI, EDDIE, (92)
EDMONDS, ROBERT L., 704
EID, S. A., 318
ELGIN, J. H., Jr., 830
ELLETT, C. WAYNE, (995)
ELLIOTT, EDWARD, S., (371)
EMGE, ROBERT G., 1071
ENDO, R. M., (441)
ENGELHARD, ARTHUR W., 894, 939
EPSTEIN, A. H., (660), 866
ERCEGOVICH, C. D., (6)
ESLAMI, A. K., (1067)
EVANS, D. W., (830)
EYAL, Z., 983

FARRAR, L. L., (362)
FARRELL, K. M., (551)
FEDER, W. A., (814), (832)
FELDMESSER, JULIUS, 476
FIELDS, W. G., (275)
FINNEY, ROBERT E., (827)
FISHER, J. R., (467)
FISHER, V. J., (65), (237)
FLEMING, A. A., 575
FORD, RICHARD E., 1014
FORER, L. B., (759)
FORSBERG, J. L., (995)
FORTIER, FREIDA, (223)
FOSTER, H. H., 169
FOSTER, ROBERT E., II, 443
FRANK, J. A., 348
FRANK, Z. R., 600
FRANKIE, GORDON W., 992
FRANKLIN, E. W., (543)
FRASER, L. R., (80)
FREDERIKSEN, RICHARD A., (682)
FREEMAN, T. E., 1043
FRENCH, W. J., 135
FREYTAG, P. H., (652)
FRICK, E. L., 770
FRIDRICH, JAMES, (451)
FROILAND, GARY E., 737
FUNK, A., 645

GAMBRELL, C. E., Jr., (169)
GARDNER, M. W., 313, (852)
GEARY, T. F., 820
GERALDSON, C. M., (953)
GIAMALVA, M. J., (622)
GIFFORD, J. R., (154)
GILMER, R. M., 133, (1062)
GILPATRICK, J. D., 39, 451, 456
GOGGANS, J. F., (192)
GOLDEN, A. M., (476)
GOLLIFER, D. E., 597
GOODING, GUY V., Jr., 92
GOSS, ROY L., (565)
GOTO, MASAO, 490
GRACEN, V. E., 432
GRAY, F. A., (472)
GRAY, L. E., 580
GROGAN, C. O., (432)
GROSKOPP, M. D., (400)
GUDAUSKAS, ROBERT T., (614)
GUMPF, D. J., (27), (441), 869
GUPTA, V. K., (693)
GUPTON, C. L., 593

HAGEDORN, D. J., 325
HAMPTON, R. O., (131)
HANCOCK, J. G., 973
HARDING, PAUL R., Jr., 256
HARDISON, JOHN R., 55, 252
HARNISH, W. N., (721)
HARPER, C. R., (774)
HARRIS, H. B., (146)
HARRISON, A. L., 873
HART, JOHN H., 685
HARTILL, W. F. T., 708
HASAN, S., 858
HAYWARD, A. C., 446, 904
HEICHEL, G. H., 1046
HELBURG, LAWRENCE B., (912)
HELFRICH, R. M., (268)
HEPER, ERCAN, (352)
HEPTING, GEORGE H., 406
HEUBERGER, J. W., (736)

HINE, R. B., 472
HOBBS, C. D., (45)
HOCK, W. K., 855
HOFF, BETTY, (863)
HOITINK, H. A. J., 480
HOLCOMB, G. E., 69, (272)
HOLDAWAY, BOYD F., 971
HOLLIS, JOHN P., 420
HOOKER, A. L., (805)
HORN, N. L., 368, 515
HORROCKS, R. D., 11
HORVÁTH, J., 665
HOUSE, J., (935)
HOWARD, CHARLES M., 33, (894), 998
HOWELL, F. C., 987
HOWELL, R. K., 195
HUBER, D. M., 342
HUDLER, GEORGE W., (712)
HUNTER, J. E., 689
HUTCHISON, D. J., 747
HYRE, R. A., 126

IBRAHIM, I. K. A., 882
IBRAHIM, ISMAIL A., (882)
INTRIERI, CESARE, 590
ISHII, M., 331
IVEY, HENRY, (362)

JEDLINSKI, H., 230
JENKINS, P. T., (858)
JENSEN, H. J., (520)
JEWELL, F. F., 531
JOFFE, A. Z., 963
JOHNSON, DENNIS R., (1071)
JOHNSON, FOLKE, 467, 779
JOHNSON, S. R., 661
JOHNSTON, R. H., (728)
JONES, A. L., 275
JONES, JOHN PAUL, 151, (504), (695), 953
JONES, R. K., (118)
JUSKA, F. V., 227, 568

KANTZES, JAMES, (783)
KATCHO, ZUHAIR A., 824
KAVANAGH, T., 235
KEIL, H. L., (844)
KELLEY, WALTER D., (192)
KenKNIGHT, G., (276)
KHADR, A. S., 878, 1040
KING, PEGGY S., (1093)
KINGSLAND, GRAYDON C., 1087
KITSOS, G. T., (264)
KLINE, D. M., 183, 891
KLOTZ, L. J., (927)
KNAUSS, J. F., 211, 391, 394, 1074
KNUTSON, DONALD M., 719
KOMMEDAHL, THOR, (861)
KOTHEIMER, JOHN B., (1004)
KRAFT, J. M., 398, (830)
KRAUSE, R. A., (836)
KREITLOW, K. W., (227)
KROHN, R. F., (241)
KRUPINSKY, J. M., 648
KUHN, C. W., 146
KUNIMOTO, R. K., (689)
KUSS, F. R., 721

LACASSE, N. L., (6), (386), (701)
LAEMMLEN, F. F., (73)
LAIRD, E. F., Jr., 722
LAMEY, H. A., 106
LANSAC, MICHELINE, (293)
LATHAM, ARCHIE J., 176
LATORRE, BERNARDO A., 1079
LAUGHLIN, C. W., 546
LAUT, JOHN G., 912
LEAR, BERT, 375
LEATH, K. T., 75
LEE, THOMAS A., Jr., 630
LEONARD, K. J., 834
LEONE, IDA A., (85)
LEWIS, J. A., (15)
LIENK, S. E., (451)
LIM, S. M., 805
LINDBERG, G. D., (157)
LINDERMAN, R. G., 1101
LINE, ROLAND F., 636
LINN, M. B., (1054)
LINZON, S. N., 526
LITTLEFIELD, LARRY J., (737)
LOGAN, JOHN, (1047), (1049)
LONG, D. L., (410)
LOONAN, DANIEL V., (659), (660)
LOUGHEED, E. C., 543
LUCAS, L. T., 774
LUKENS, R. J., 203
LUNG, M. C. Y., (652)

MAAS, JOHN L., 296
MacKENZIE, D. R., 164
MAGASI, LASZLO P., 245
MAGIE, R. O., (939)
MAHMOOD, TARIQ, 582
MAI, W. F., 141, 434, (577)
MAIRE, R. G., (540)
MAKKOUK, K. M., 272
MALEK, R. B., (898)
MALO, S. E., (1000)
MANANDHAR, JUJU B., (594)
MANNING, W. J., 9, 405, 814, 832
MARAMOROSCH, KARL, (643)
MARCO, S., 1035
MARENAUD, C., (293)
MARLATT, ROBERT B., 471, 1091
MARSH, PAUL B., (215)
MARTIN, W. J., 58, 219
MARTINSON, C. A., (29)
MASON, LEE, 485
MASSOUD, S. I., (882)
MATEJKA, JOSEPH C., (773)
MATHRE, D. E., 728
MATOBA, FRANK, (1051)

MATTHEWS, FRED R., 931
McCAIN, A. H., 241
McCORNACK, A. A., (909)
McCOY, R. E., 1019
McGRATH, W. THOMAS, 99
McILVEEN, W. D., (526)
McMILLAN, R. T., Jr., 759, 967, 969
McNEAL, F. H., (761)
McNEILL, M. J., 104
McVEY, D. V., (410), (554), (1038)
MEADE, NYMPHES, (817)
MEISTER, H., (935)
MERRILL, W., 525, 1058
MEW, I-PIN CHANG, 861
MILHOLLAND, R. D., 118
MILLER, D. L., (498), (939)
MILLER, DOUGLAS R., 843
MILLER, L. A., (162)
MILLER, P. M., 255, 612, 1102
MILLIKAN, D. F., (462)
MINK, G. I., 93, (198)
MIXON, AUBREY C., (415)
MOORE, L. D., (3)
MOORE, RICHARD, (1102)
MOREY, D. D., (534)
MORGAN, PAUL D., 407
MORROW, RONALD A., (912)
MORSE, FRANK S., (99)
MUKHOPADHYAY, A. N., 776
MULANAX, M. W., (342)
MULLANY, R., (203)
MULLER, G. J., 1054
MUSE, R. R., 1083

NAGARAJAN, K., (740)
NAUER, E. M., (976)
NEELY, DAN, 354, 460, 667
NELSON, J., (935)
NELSON, L. R., 534
NELSON, M. R., 731
NELSON, P. E., (836)
NELSON, R. R., (59), (164), 767, (889)
NEMEC, S., 345
NEWTON, H. CALVIN, 798, 875
NICHOLLS, THOMAS H., 712
NICHOLS, LESTER P., 252, (525), 759
NIGH, EDWARD L., Jr., (507), 914
NISHIJIMA, W. T., (73)
NORTON, R. A., (792)
NOVACKY, ANTON, 765

O'BANNON, J. H., (747)
OGBUJI, R. O., 520
OLTHOF, TH. H. A., (417)
ORMROD, D. J., 53
ORR, CALVIN, C., 1065
OSBORN, JOHN, (223)
OTEIFA, B. A., (1040)
OTTA, J. D., 572
OU, S. H., (960)
OVERMAN, A. J., (953)

PAIR, J. C., (957)
PAISLEY, R. B., 736
PALMER, L. T., (740)
PAPAVIZAS, G. C., 15
PAPIA, P. M., (9)
PARKER, K. G., (141)
PARMETER, J. R., (992)
PARTRIDGE, A. D., 498, 939
PATEL, B. N., (210)
PATEL, M. V., (210)
PAULUS, A. O., 935
PAWUK, WILLIAM H., 944
PAYNE, K. T., (32)
PEARSON, R. G., (526)
PEET, CLYDE E., (104)
PELLISSIER, M., 6
PERKINS, I., (814), (832)
PÉROMBELON, M. C. M., 552
PERRY, D., (241)
PERRY, R. S., 995
PERRY, S. K., (378)
PETERSON, ROGER S., 474, 896
PETROVICH, M. S., (808)
PEVERLY, G., (112)
PFLEGER, FRANCIS L., 180
PHELPS, RALPH H., (817)
PIRONE, T. P., 652
PLAVŠIĆ-BANJAC, BILJANA, 643
POEHLMAN, J. M., (167)
POLITIS, DEMETRIOS J., 1026
PONELERT, C. G., (652)
POTTER, J. W., 417
POVISH, W. R., (336)
POWELSON, R. L., 178
PRESCOTT, J. M., (740)
PREY, ALLEN J., (99)
PROVVIDENTI, R., 281, (285), 548

RADEWALD, J. D., 540
RAFII, Z., (206)
RAJAGOPALAN, B., 323
RAMEY, HARMON H., Jr., (215)
RAND, R. E., (325)
RANDHAWA, G. S., (990)
RAO, V. RAMA, (990)
RASKI, D. J., (1028), 1031
RAYCHAUDHURI, S. P., (424)
REEDER, W. F., (751)
RHODES, W. H., (169)
RICH, AVERY E., (1004)
RICH, SAUL, (1102)
RICHARDSON, M. J., 803
RIEDEL, R. M., (329)
ROANE, C. W., (183)
ROBERTS, JOHN J., (827)
RODRIGUEZ-KABANA, R., 362, 1093
ROELFS, A. P., 410, 554, 608, 1038
ROGERS, KENNETH M., 415
ROHDE, C. R., (178)
ROISTACHER, C. N., (976)
ROTHMAN, P. G., (608)

ROWAN, SAMUEL J., 558, 628, (931)
ROWELL, J. B., (410)
RUEHLE, JOHN L., 691
RUSH, M. C., 154, (157), 622
RUSSELL, H., (15)
RYUGO, KAY, (590)

SAAD, S. M., (325), 744
SAETTLER, A. W., 378, (714)
SAHARAN, G. S., 693
SALADINI, J. L., 822, 885
SALEM, A. A., (1040)
SALLY, B. K., (188)
SANDHU, R. S., (601)
SCHALLER, C. C., 276
SCHEFFER, R. P., (247)
SCHENCK, N. C., 276
SCHIEBER, EUGENIO, 753
SCHMIDT, R. A., 811
SCHMITT, R. V., (1031)
SCHMITTHENNER, A. F., 536
SCHOENEWEISS, D. F., 238
SCHONHORST, M. H., (472)
SCHREIBER, L. R., (855)
SCHROEDER, W. T., (281), 285, (548)
SCHUSTER, M. L., 139, 863
SCOTT, L. E., 783
SECHLER, DALE T., (167)
SEEM, R. C., 386
SELF, R. L., (42), (919)
SELVARAJ, Y., (990)
SEQUEIRA, LUIS, (798), (875)
SETH, M. L., 424
SEYMOUR, C. P., (511), (811)
SHANER, GREGORY, 358, 827
SHARP, E. L., (188), 761
SHEIDOW, N. W., (417)
SHER, S. A., (540)
SHIBUYA, F., (540), (935)
SHIGO, ALEX L., (406)
SHORTLE, WALTER C., 1004
SIMBWA-BUNNYA, M., 901
SIMPSON, MARION E., 215
SINCLAIR, J. B., (1054)
SINCLAIR, W. A., 159
SINGH, D. V., (424)
SINGH, N. D., 339, 551, 1059
SINGH, RAMA S., 115
SKELLY, J. M., 3
SMITH, C. A., (39), (451)
SMITH, D. H., (290), 796
SMITH, R. B., (543)
SMITH, RICHARD S., Jr., 102, 241
SMITH, R. T., (648)
SMITH, S. H., (785)
SMITH, W. L., Jr., (296)
SMOLIK, J. D., 898
SNOW, G. A., (531)
SNYDER, W. C., (493)
SONODA, RONALD M., 840
SOOST, R. K., (927)
SOULEN, D. M. 785
SPALDING, D. H., 751
SPIERS, JAMES M., (917)
SPRINGER, J. K., 162
SPURR, HARVEY W., Jr., 338
SRAGO, M., (241)
SRIDHAR, R., 960
STAMBAUGH, W. J., (987)
STANGHELLINI, M. E., 507
STATLER, GLEN D., 77, 788
STEWART, JOSEPH K., 1051
STIPES, R. JAY, 108, 604
STONE, L. L., (3)
STOUFFER, R. F., (785)
STOVER, R. H., 1000
SU, SHU-HUA, (682)
SUBIRATS, F. J., 42, 919
SUNDARAM, N. V., 740
SZKOLNIK, M., (456)

TARJAN, A. C., 186, 626
TEWARI, I., (633), (676)
THAKUR, R. P., (776)
THANASSOULOPOULOS, C. C., 264
THAYER, P. L., 45
THOMAS, C. A., (195)
THOMPSON, S. S., 290
TOLER, ROBERT W., (69), (630), 658
TOMPKINS, C. M., 415, 493
TROUTMAN, JOSEPH L., 773
TRUJILLO, GUSTAVO E., 714
TSAO, PETER H., (971)
TULEEN, DARLENE M., 682
TUNG, GABRIELLE, (767)
TURNER, M. T., 29
TURNER, N. C., (1046)

UYEMOTO, J. K., (133), 1062

VAARTNOU, H., 633, 676
Van ARSDEL, E. P., 300
van der MESPEL, G. J., (850)
van der ZWET, T., 844
VARDARO, P. M., (405)
VARGAS, J. M., Jr., 32, 334, (546)
VARNEY, E. H., (162)
VAUGHAN, EDWARD K., (180)
VAZIRI, A., (1067)
von UEXKULL, HELMUT R., (643)

WADE, E. K., (744)
WAGNER, LARRY J., (139)
WALLIN, JACK R., 659, 660, (866)
WALTON, G. S., (1046)
WARREN, H. L., 268
WATERWORTH, HOWARD E., 336, 923
WATERWORTH, Y. L., (80)
WATTERSON, J. C., 949
WEATHERS, L. G., 27, 441, (869)
WEERAPAT, PRAPHASE, 167
WEHUNT, E. J., 305

WEIDENSAUL, T. CRAIG, 701, 942
WEINKE, K. E., (604)
WESTER, HORACE V., 725
WESTER, ROBERT E., 65, 237
WHEELER, HARRY, (1026)
WHEELER, R. E., (731)
WHITAKER, T. W., (935)
WHITE, DONALD G., 995
WHITESIDE, J. O., 671
WILCOXSON, R. D., 286
WILFRET, GARY J., (894)
WILLARD, J. R., 808
WILLIAMS, P. H., (949)
WILLIAMS, R. J., (106)
WILLIAMS, WAYNE T., 223
WILLIS, W. G., (957)
WILSON, D. R., (348)
WILSON, K. I., (323)
WITCHER, WESLEY, (310)
WOLTZ, S. S., (151), (695)
WOOD, FRANCIS A., (942), (944)

YARWOOD, C. E., (313), 852, 1085
YORK, P. A., (261)

ZAGAJA, S. W., 462
ZETTLER, F. W., (822), (885)
ZILLINSKY, F. J., (803)
ZIMMER, D. E., 428
ZITTER, THOMAS A., 586
ZUBER, M. S., (11), (485)

SUBJECT INDEX

Abies grandis (grand fir): significant bark beetle-root rot association in grand fir in Idaho, 498
Abscission: of foliage of Ficus elastica by Cercospora disease, 1091
Acacia koa (koa): collar rot of koa, caused by Calonectria crotalariae, 73
Adjuvants (see also Antibiotics, Chemicals, Fungicides, and Nematicides): Folicote as an antitranspirant reduced ozone injury on bean plants, 6; pinolene (Nu-Film 17), used with benomyl gave excellent control of strawberry leaf blight, 23
Aecidium garkeanum: 1047
Africa: 105; 318; 678; 755; 901; 1011; 1047
Agave: Phytophthora isolates from zebra disease tested for pathogenicity, 678
Ageratum: host of tomato spotted wilt virus in nursery stock in California; 869
Agrobacterium tumefaciens: 108
Air pollution: oxides of nitrogen and sulfur dioxide caused severe reduction in growth and tip burn in eastern white pine seedlings, 3; chrysanthemum varieties are apparently not sensitive to ozone damage under field conditions, 85; sensitive cultivars of safflower are suggested as useful biological indicators for phytotoxicants, 195; results of the 1969 statewide survey of air pollution damage to vegetation in Pennsylvania, 701
Ajuga reptans: first report of alfalfa mosaic virus in Ajuga reptans, from New York, 285
Alabama: 42; 415; 614; 919
Alaska: 618; 645
Alberta: 633; 676
Alfalfa: control of leaf diseases with fungicides, and yield increase, 286; Phytophthora root rot caused by Phytophthora megasperma in Arizona, 472; found in Washington, 830
Allium cepa (see Onion)
Alternaria sp.: 328; 721
--- alternata (Alternaria tenuis): 633; 676; 776
--- brassicae: 633
--- citri: 909
--- dianthi: 9; 728
--- raphani: 633
--- solani: 308
--- tenuissima: 118
Altitudinal zones of parasitism: for three species of dwarf mistletoe and their pine hosts in Colorado, 223
Anatomical mechanism: for rust resistance, in slash pine, 531
Announcement: instructions to authors submitting manuscripts to the Plant Disease Reporter, 368; 464
Announcement: address change for correspondence with the Plant Disease Reporter, 1016; 1104
Anthracnose: of cucurbits, Colletotrichum orbiculare isolates from Oklahoma show differential pathogenicity, 35; of sycamore, control of twig blight stage with fungicide phenylmercuritriethanolammonium lactate, 354; of strawberry, control of crown rot phase through sanitation, breeding for resistance, and benomyl fungicidal control, in Louisiana, 515; of mango, hot water treatment controlled anthracnose as did benomyl and thiabendazole but caused scald injury, 751; the perfect stage of Colletotrichum graminicola, 1026; caused dieback of regrowth on defoliated oak and maple, in Connecticut, 1046; of kenaf, caused by Colletotrichum hibisci, a new race appears to be present in Zambia, 1049

Anthurium andraeanum: cv. Kansako Red found infected with Xanthomonas dieffenbachiae in Hawaii, a first record of this pathogen on anthurium outside of Brazil, 904
Antibiotics: cycloheximide, an isolate of Cochliobolus carbonum with single gene fungicide tolerance to cycloheximide was tested under field conditions, 164
---: streptomycin, some isolates of Xanthomonas dieffenbachiae showed resistance to streptomycin spray on Philodendron oxycardium, 394
---: tetracycline, remission of lethal yellowing in coconut palm treated with tetracycline antibiotics, 1019
Aphelenchoides besseyi: 471
Apple: Hormiscium sp. infected 'Jonathan' apples in cold storage in Michigan, 275; excellent control of scab, powdery mildew, rusts and European red mite on apple with new piperazine derivative fungicide, 451; stem-cavity browning of 'McIntosh' apples, 543; preplant fumigation with three nematicides resulted in increased growth of trees and reduced populations of Pratylenchus penetrans, 577; apple leaf spot, an Alternaria sp. was found associated with lesions, 721; powdery mildew, fall applications of pruning agents to mildewed shoots yielded almost total eradication of the pathogen, 770
---: crab, scab infections on additional species and varieties of flowering crab apples, 252
---: scab: a postharvest pre-leaf-fall spray of apple trees with benomyl reduced primary scab infection the following spring in Delaware, 736; controlled in Chile with late season applications of benomyl + surfactant F, 1079
Arachis hypogaea (see Peanut)
Arceuthobium americanum: 223
--- cyanocarpum: 223
--- vaginatum: 223
Arizona: 472; 507; 731; 773; 914
Arkansas: 270; 355; 658
Armillaria mellea: 406; 498
Artificial infestation: of large field plots cropped to peas with Fusarium solani f. sp. pisi, 398
Ascension Island: 869
Ascochyta coffeae: 753
--- gossypii: 328
Aspergillus flavus: 215
--- glaucus: 173
--- niger: 64
--- restrictus: 173
Assay: local-lesion, 'Monroe' bean produced consistent formation of local lesions when rub-inoculated with bean common mosaic virus, 714
Australia: 53; 80; 290; 645; 858
Avena (see Oat)
Avocado: the most common Fusaria associated with avocado fruit in Israel are F. moniliforme and F. equiseti, 963
Azalea: control of stunt nematode with DBCP, 540; occurrence of azalea petal blight in Connecticut, 1101

Bacillus subtilis: 861
Bacterial diseases: of bean in Michigan, 378
---: bacterial blight: of carrot in Wisconsin, 744; of cotton, infection of cotton cotyledons with Xanthomonas malvacearum causes darkening of internal pigment glands, 765
---: bacterial brown spot of bean, caused by Pseudomonas syringae, cultivars and plant introductions tested for tolerance, 325
---: bacterial canker: of tomatoes, controlled in water culture with formalin, 139; bacterial canker and bacterial soft rot were the tomato diseases causing the greatest yield losses in Montserrat, West Indies, 817
---: bacterial leaf spot: leaf spot and tip burn of Philodendron, isolates of Xanthomonas dieffenbachiae showed resistance to streptomycin sprays, 394; of clover, caused by Pseudomonas stizolobii in Hawaii, 446; of onion, in Japan, caused by Pseudomonas syringae, 490; of tomato, relative susceptibility of some tomato genotypes to Xanthomonas vesicatoria, 504; Xanthomonas dieffenbachiae found on anthurium leaves in Hawaii, 904
---: bacterial stem rot of poinsettia, 480
---: bacterial wilt of cucumber, movement of Erwinia tracheiphila, 949
---: fire blight of pear, prevalence of blight infection in 'Magness' trees, 844
---: Nebraska leaf freckles and wilt bacterium of corn, 863
Banana: Fusarium moniliforme is the cause of "black heart" of banana, in Israel, 963; occurrence of Fusarial wilt in normally resistant 'Dwarf Cavendish' banana in the Canary Islands, 1000
Bajra streak: a leafhopper-borne cereal virus on pearl millet in India, 424
Bark split disease: of prune, possibly caused by a type of apple chlorotic leaf spot virus, 293
Barley: benomyl as a preplanting soil drench gave good control of cyst production of Heterodera avenae on barley, 261
---: Helminthosporium stripe: several fungicides as seed treatments reduced incidence of disease, 183; 27 spring barley cultivars were immune and 13 showed symptoms in

less than 1% of seedlings tested for resistance to Helminthosporium stripe, 891
Basidiomycete: a sterile basidiomycete apparently caused a spotting and discoloration on dormant bermudagrass in Arkansas, 355
Bean: evaluation of varieties and selections of lima beans for resistance to Rhizoctonia solani, 268; bean common mosaic virus, 'Monroe' has use as a local-lesion assay host for BCMV because lesion production is uniform and consistent, 714; new race of bean rust found on 'Arthur' pole beans in Florida, 759; zinc + maneb was the most effective of four fungicides for control of bean diseases in Kawanda, Uganda, 901; new disease, wet-rot of pole bean, in Florida, 967
---: bacterial diseases: 180 commercial cultivars and 503 bean plant introductions tested for reaction to bacterial brown spot disease, 325; seed-transmitted bacterial diseases in Michigan, 378
---: downy mildew: lima bean breeding lines carry multiple resistance to downy mildew, 65; occurrence of downy mildew in lima beans in New Jersey in 1971, 237
---: ozone injury: wax emulsion sprays on leaves of 'Pinto III' bean gave protection against ozone injury, 6; ozone injury to 'Pinto III' seedlings suppressed with triarimol and EL-279, 386
Beech: late-spring leaf scorch is physiological in nature, 526
Beet (red table): under greenhouse conditions growth of beets was markedly reduced when inoculum level of Heterodera schachtii reached 2443, 434
Beetle-root rot relationship: in Idaho conifers, 498
Beggarweed (Desmodium tortuosum): potassium azide broadcast post-emergence to peanuts completely controlled beggarweed in Florida test, 362
Belonolaimus longicaudatus: 957
Bifusella linearis: 102
--- saccata: 102
Bioassay: leaf whorl bioassay very reliable for determining susceptibility of corn lines to Helminthosporium maydis, 29; of fungicides for control of Phytophthora blight of macadamia, 689; root extracts of sugarbeet seedlings treated with triphenyltin chloride inhibited Alternaria tenuis on PDA, 776; inhibition of Penicillium atrovenetum and Rhizoctonia solani by extracts of tomato seedlings exposed to three concentrations of thiabendazole and ETMT, respectively, 1054
Biological control: Puccinia chondrillina will infect skeleton weed over a wide temperature range and will remain host-specific under many different climatic conditions, 858
Biotypes: of Meloidogyne hapla in the Pacific Northwest, 520
Birdsfoot trefoil (Lotus corniculatus): Stemphylium loti was most important pathogen on southern slope exposure in West Virginia, 731
Black shank: of tobacco, race 1 of Phytophthora parasitica var. nicotianae is present in east Tennessee, 593
Blast disease: of ryegrass, reported for the first time from Louisiana in 1971, 157; an epiphytotic condition on ryegrass in Mississippi in 1971, 210; of finger millet in India, 740; lesion development on a rice cultivar by different races of the blast pathogen, 960
Blight: Ascochyta blight, estimated cotton loss in U.S. in 1971, 328; aerial blight of Carissa grandiflora by Rhizoctonia solani, 511; Arizona cypress blight, control in nursery plantings with fungicides, 558
---: bacterial: estimated cotton loss from Xanthomonas malvacearum by State in 1971, 328; survey of internally-borne bacterial diseases in Michigan navy beans, 378; of carrot, found for first time in Wisconsin, 744; cotton, internal pigment glands darken when Xanthomonas malvacearum infects the cotton cotyledons, 765
---: fire: of pear, incidence in trunks of 'Magness' pear trees, 844
---: foliar blight of Dionaea muscipula, caused by Colletotrichum gloeosporioides, 391
---: leaf: of strawberry, Dendrophoma obscurans controlled with benomyl, 23; of centipedegrass, 42; of coffee, caused by Ascochyta coffeae, 753; of wheat, caused by Fusarium nivale in Scotland and on triticale in Mexico, 803; of tomato, caused by Phoma destructiva, 869
---:---: southern corn leaf blight (SCLB): relationship between stalk lodging and corn leaf blight, 11; potential for spread and increase, 104; varying levels of susceptibility and resistance to H. maydis in corn inbreds and hybrids of different cytoplasms and genotypes, 112; effect of shade and wet soil on incidence, 203; 1971 results of interregional and regional corn leaf blight ratings, 485; does not appear to endanger sorghum production at present time, 682; on sweet corn with T cytoplasm in Florida, 969; data indicate H. maydis can overwinter on crop debris on the soil surface and initiate the primary disease cycle in the spring in western South Carolina, 1087

---: mukwa blight in Central Africa, 820
---: needle, brown spot needle blight of Scotch pine Christmas trees in Wisconsin, 99
---: petal blight of azalea reported from Connecticut, 1101
---: Pythium blight of ryegrass, ETMT seed treatments gave effective control, chloroneb did not, 1043
---: seedling blight of wheat, caused by Helminthosporium sorokinianum or Fusarium roseum or a combination of the two, 788
---: stem, of Vinca minor, caused by Phoma exigua var. exigua, 238
---: stem-end blight of pecan, reduced by benomyl and TPTH in Louisiana, 276
---: Sirococcus tip blight of Jeffrey pine seedlings, controlled with fungicides, 241
Blueberry: postharvest decay in North Carolina berries from Alternaria tenuissima, 118
Blue mold: of citrus, control of Penicillium mold decay and sporulation by thiabendazole is dependent upon the nature of the strains involved, 256
Botryodiplodia hypodermia: 572
--- theobromae: 990
Botryosphaeria dothidea: 108
Botrytis cinerea: 126; 296; 814
British Columbia: 53; 645
British Solomon Islands Protectorate: 597
Brown spot (Scirrhia acicola): of red pine, a new host of the pathogen, 712
Brush burning: on preplant sites of sour cherry increased growth and yield over untreated trees, in New York, 141
Brussels sprouts (see Crucifers)
Burned sites (see Fire damage)

Cabbage (see Crucifers)
Cacao (cocoa): isolation of Phytophthora palmivora from cocoa bark cankers, 71; 'Amelonado' seedlings used to study an isolate of cocoa swollen shoot virus from Nigeria that differs from other isolates, 755
Cadang-cadang: a disease of coconut palm from the Philippines, 643
Caliciopsis pinea: 108
California: 27; 102; 313; 375; 441; 493; 722; 843; 852; 869; 935; 973; 1031
Calonectria crotalariae: 73
Camellia: a flower rot disease of camellias in Mobile, Alabama is thought to be caused by a mixed culture of Penicillium implicatum and P. rugulosum, 919
Canada: 53; 417; 526; 633; 645; 676
Canary Islands: 1000
Canker: cocoa bark cankers used for isolation of disease pathogen, Phytophthora palmivora, 71; orange hobnail, bark slough canker, coal canker, Fusarium and Polyporus cankers found on oaks in Texas, 300; Siberian elm canker, caused by Botryodiplodia hypodermia, in South Dakota, 572; of Monterey cypress, and the probable role of the moth, Laspeyresia cupressana, as an effective disseminator of Coryneum cardinale spores that cause serious canker, 992
---: bacterial, of tomatoes, grown hydroponically, 2% formalin to flush gravel beds controlled the disease, 139
---: Fusarium: of yellow poplar in South Carolina, 310; of hybrid poplars in Ohio, 523; of maple, precipitation alone apparently does not have significant influence on frequency of cankers, 942; of maple, influence of sugar maple stem tissue extracts on spore germination of the fungus, 944
Cantaloup (see Cucurbits)
Capsella bursa-pastoris: susceptible to isolates of beet western yellows mosaic virus, 198
Carissa grandiflora: serious blight disease found to be caused by Rhizoctonia solani, 511
Carnation: benomyl-treated plants showed more Alternaria leaf spot than untreated control plants, 9
Carrot: bacterial blight of carrot, caused by Xanthomonas carotae found for first time in Wisconsin in 1971, 744
Carya illinoensis (see Pecan)
Cauliflower (see Crucifers)
Cedar: eastern red, nursery seedlings found infected by Curvularia intermedius in Alabama forest nursery, 192
Celery: nematode and soil fungi control in celery seedbeds on muck soil by flooding and drying, or with fumigants, 661
Cephalosporium sp.: 108
--- diospyri: 916
--- gregatum: 580
Ceratocystis fimbriata: 219
--- ulmi: 460; 604; 667; 685; 725; 912
Cercospora sp.: 328; 1091
--- arachidicola: 601; 1099
--- medicaginis: 286
--- personata: 601; 873; 1099
--- sequoiae: 558
Cercosporella herpotrichoides: 20; 178; 342; 594
Chaetomium globosum: 861
Chemicals (see also Adjuvants, Antibiotics, Fungicides, and Nematicides): broad spectrum soil fumigants tested for their effect on raspberry root rot control, 467; effects of P, S, N, lime, chlordane and fungicides on Ophiobolus patch on 'Highland' bentgrass putting green turf, 565

---: CO_2, injury in lettuce, 1051
---: 1-decanol, a contact suckering agent applied with triphenyltin hydroxide fungicide to tobacco caused leaf injury, 338
---: dimethyl sulfoxide may influence the translocation of alfalfa mosaic and tobacco ringspot viruses, 1014
---: K_2HPO_4 + bentonite added to Chenopodium amaranticolor virus inoculum increased infectivity of inoculum by 7- to 1200-fold, 1085
---: insecticide, PP-211 (O-[2-(diethylamino)-6-methyl-4-pyrimidinyl]O,O-diethyl phosphorothioate), a candidate insecticide improved effectiveness of four seed-treatment fungicides on rice, 154
---: lime, lime in soil mix was apparently the cause of poinsettia wilt, 415
---: sulfur dioxide: in combination with oxides of nitrogen caused damage to eastern white pine, 3; not damaging to chrysanthemum varieties, 85
---: wax, wax emulsion sprays protected bean plants from ozone injury, 6
Chenopodium amaranticolor: cucumber mosaic virus buffering affects response in C. amaranticolor, 443; virus transmission of cucumber mosaic from C. amaranticolor enhanced with additions of K_2HPO_4 and bentonite, 1085
Cherry (see also Prunus): preplant nematicide, preplant nematicide + plant nutrients, brush burning on planting sites, and complete nutrition treatments each increased growth and yield of trees over untreated trees, 141; stem pitting of sour cherry was found in Poland, 462
Chestnut (Castanea crenata): better penetration and translocation of a fungicide after foliar treatment of seedlings with growth retardants, 590
Chinese gooseberry (Actinidia chinensis): control of root-knot nematodes on 2nd-year cuttings with V-C 9-104 or cynem, 850
Chlamydospores: of Phytophthora parasitica remained viable for long periods when stored in deionized distilled water at 9°C, 971
Choanephora cucurbitarum: 967
Chrysanthemum: 16 varieties tested for response to sulfur dioxide and ozone showed little sensitivity, 85; leaf rust, caused by Puccinia chrysanthemi, in Florida, 939
Cicadulina mbila: 424
Citrus: a disease that causes sudden wilting and death of trees has been found in New South Wales, Australia, 80; a new citrus disease complex found in Tanzania, 318; hybrids of Citrus x Poncirus were tested for resistance to Phytophthora parasitica and P. citrophthora, 927
---: nematodes: Kallstroemia maxima, found to be a new host of the burrowing nematode and often reestablishes itself in the chemical barriers maintained to contain the nematode in presently infested areas of Florida, 85; extraction of citrus nematode from infested roots, 186; paste application of BAY 68138 to stem of Pratylenchus-infested citrus seedlings significantly reduced numbers of nematodes in roots of rough lemon, 626; 26 Citrus hybrids and varieties and 13 orange subfamily relatives evaluated for resistance to the citrus nematode, 747
---: fruit decay: Citrus aurantifolia (lime), Aspergillus rot found on fruit in local vegetable markets in India, for the first time, 64; lemon, strains of citrus blue mold and green mold pathogens show differential sensitivity to thiabendazole, 256; irradiation controlled sour rot decay on lemon, but caused fruit injury, 582; greasy spot rind blotch is name proposed to cover the disease syndrome caused on citrus by Mycosphaerella citri that makes rind blemishes including "pink pitting," 671; decay of citrus caused by Alternaria citri in Florida, 909; Fusarium moniliforme was most common Fusarium on grapefruit fruit in Israel, F. solani was more prevalent on oranges, 963
---: virus: vein-enation virus on Satsuma mandarins in Turkey, 352; concave gum, infectious variegation, psorosis-A, tristeza, seedling yellows-tristeza, and vein enation viruses of citrus were inactivated by 2- to 3-months' hot air treatment at alternating temperatures of 40°C for 16 hr/day and 30°C for 8 hr/night, 976
Cladosporium sp.: 990
--- epiphyllum: 108
Claviceps sp.: 740
Clitocybe tabescens: 811
Clonal differences: in elms, based on physiological vigor for crown resistance to Dutch elm disease, 725; in the incidence of surface rot on sweetpotato, 783
Clover: a bacterial leaf spot on clover (Trifolium spp.) found in Hawaii was caused by Pseudomonas stizolobii, 446; Ladino, several viruses not reported previously from white clover from North Carolina were isolated and identified, 774
Clubroot: a combined application of nematicide and PCNB was necessary to control sugarbeet nematode and the clubroot organism on Brussels sprouts, 375
Cochliobolus carbonum: 164
Cocoa (Theobroma cacao): technique developed for isolation of Phytophthora

palmivora from cocoa bark cankers, 71; peculiarity of an isolate of cocoa shoot virus from Nigeria, 755
Coconut palm: preliminary observation of cadang-cadang-diseased coconut palm leaves from the Philippines by electron microscopy, 643; direct injection of oxytetracycline-HCl or tetracycline-HCl into lethal yellowing-affected coconut palms stopped symptom development and healthy new growth appeared, in Florida, 1019
Coffee: a leaf blight on coffee, caused by Ascochyta coffeae, has been found in the highlands of Guatemala, 753
Coleosporium crowellii: 474
Collection efficiency: of rotorod samplers for sampling fungus spores in the atmosphere, 704
Colletotrichum sp.: 894
--- coccodes: 321
--- fragariae: 515
--- gloeosporioides: 391; 751
--- graminicola: 1026
--- hibisci: 1049
--- orbiculare: 35
Colocasia esculenta: two virus diseases, Alomae and Bobone, are described for the first time from the British Solomon Islands that infect taro or dasheen (Colocasia esculenta) a major subsistence crop of the Islands, 597; a new host of Meloidogyne incognita in Trinidad, 618
Colorado: 223; 912
Conifers (see also Species names): bark beetles and root rots show relationship in Douglas-fir, grand fir, and ponderosa pine in Idaho, 498
Connecticut: 1046; 1101; 1102
Contaminants: Helminthosporium spp. can develop readily as contaminants on leaf specimens incubated in moist chambers, 834
Coprinus micaceus: 80
Corms: of Xanthosoma spp., dasheen mosaic virus not inactivated in corms with hot water treatments below 65°C, above this, corms ceased to germinate, 320; corm disease, of Gladiolus, caused by Curvularia trifolii sp. gladioli, 995
Corn: relationship between stalk lodging and corn leaf blight, 11; monogenic tolerance of Cochliobolus carbonum isolates to cadmium succinate and cycloheximide fungicides tested under field conditions, 164; Phaeosphaeria herpotricha found on southern corn leaf blight-infected plants rather than Cochliobolus heterostrophus as first reported, in Florida, 276; male cytoplasmic effect on reaction of corn to diseases, 575; a disease of corn that causes stunting, has viruslike particles, and is leafhopper-transmitted, has been found in Kentucky, 652; stalk rot fungi in Pennsylvania in 1970 and 1971, 836; recovery of fungi from surface-disinfected kernels where fungi were known to be present was reduced by application to the kernels before plating of storage fungi and soil organisms, 861
---: leaf freckles and wilt (also called Goss's wilt), most of the 10-day-old seedlings of 100 corn inbreds inoculated with two isolates of the wilt bacterium were quite susceptible, 863
---: popcorn, storage tests indicate that a combination of moisture content below 14.0% and temp of 10-20°C should be suitable for long-time storage of popcorn without loss in quality from invasion by storage fungi, 1097
---: southern corn leaf blight (SCLB): 25 sources of cytoplasmic male sterility tested by leaf whorl bioassy with toxin of Helminthosporium maydis race T, 29; the potential for spread and increase of SCLB, 104; reaction of corn inbreds and hybrids with different cytoplasms and genotypes to H. maydis, race T, 112; corn foliage in morning shade and on wet soil has a greater incidence of SCLB than that growing in full sun or on dry ground, 203; reactions of corn seedlings with nonsterile, Texas male-sterile, and restored Texas male-sterile cytoplasms to H. maydis toxin, 432; leaf blight ratings of corn in several cytoplasms from the interregional and regional leaf blight rating tests for 1971, 485; incidence of H. maydis race T in Iowa in 1971, 660; infections of corn hybrid with race T of H. maydis increased with increasing temperatures or dew periods, 767; SCLB development in Iowa monitored by network of uniform blight indicator plots across the State, 866; a dramatic decrease in populations of race T is predicted in areas that grow predominantly normal cytoplasmic hybrids in which populations of race O occur, 889; SCLB found on T cytoplasm sweet corn in Florida, 969
Correction: (Vol. 55, No. 11, Page 960), 96
Correction: (Vol. 55, No. 11, Page 996), 96
Correction: (Vol. 55, No. 11, Page 1037), 192
Correction: (Vol. 56, No. 1, Page 29), 920
Correction: (Vol. 56, No. 2, Page 109), 560
Correction: (Vol. 56, No. 3, Page 206), 367
Correction: (Vol. 56, No. 3, Page 210), 560
Correction: (Vol. 56, No. 8, Page 705), 1104
Corynebacterium sp.: 863
--- flaccumfaciens var. aurantiacum: 378
--- michiganense: 139

Corynespora cassiicola: 243
Coryneum cardinale: 992
Cotton: six surveys suggest that Aspergillus flavus boll rot is restricted geographically within the U. S., 215; reduction in yield of cotton caused by parasitic diseases in the U. S. in 1971, 328; Longidorus siddiqii shown to have pathogenic capacity to infect Egyptian cotton, 699; influence of Xanthomonas malvacearum infection on cotton pigment glands, 765; capacity of Pythium splendens to cause preemergence damping-off of cotton seedlings was compared with other Pythium spp., 973; varietal susceptibility and significance of the reniform nematode in Fusarium wilt of cotton, 1040; DBCP fumigation for narrow-row cotton on the Texas High Plains, 1065
Cowcockle (Saponaria vaccaria): Alternaria dianthi is cause of leaf spot on this starch-producing plant, 728
Crambe (see Crucifers)
Creeping bugleweed (see Ajuga reptans)
Criconemoides onoensis: 420
--- ornatus: 476
Cristulariella sp.: 108
--- pyramidalis: 135; 176; 796; 917
Crocus (Colchicum sp.): tobacco ringspot virus was isolated from 'The Giant' crocus in Maryland, 336
Cronartium fusiforme: 531; 628; 931
Crownvetch (Coronilla varia): Sclerotinia trifoliorum caused severe crown and root rot on northern exposures but was never found on south-facing exposures, 371
Crucifers: Brussels sprouts, control of sugarbeet nematode with several low vapor pressure nematicides in California, 375
---: cabbage, greenhouse study on relationship between population levels of Heterodera schachtii and root and top weights of red table beet and cabbage, 434
---: cauliflower: turnip mosaic virus was isolated from 'Green Ball' cauliflower, 441
---: Crambe, several Crambe species were tested for their reaction to several plant viruses, 665
---: rape (Brassica campestris): Alternaria alternata, A. brassicae, and A. raphani were found to be parasitic on Polish-type rape in Alberta, Canada, 633; pathogenicity of A. alternata on Polish-type rape in Alberta, Canada, 676
Cucumber (see Cucurbits)
Cucurbits: six Oklahoma isolates of Colletotrichum orbiculare showed differential pathogenicity to various cultivars of watermelon, cucumber, squash, and cantaloup, 35; Macrophomina phaseoli isolated from Lagenaria vulgaris, Cucurbita pepo, Luffa acutangula, Cucumis melo, and Citrullus lanatus, in India, 115; importance of Momordica charantia as a watermelon virus 1 source for cucurbits in Palm Beach County, Florida, 563
---: cantaloup, triarimol and PP 675 gave excellent control of cantaloup powdery mildew, 935
---: cucumber: Phomopsis root rot found in British Columbia greenhouse, first known occurrence in North America, 53; triarimol and wettable sulfur evaluated under field conditions for control of powdery mildew on cucumber in Iran, 206; Erwinia tracheiphila movement and multiplication in resistant and susceptible cucurbits, 949
---: squash: powdery mildew controlled with triarimol in Florida and Texas, 45; watermelon mosaic virus considered the most serious disease of summer squash in Georgia, 147; squash and cucumber, PP 675, a systemic fungicide, was more effective against powdery mildew than two standard fungicides Karathane or sulfur, 878
---: watermelon: identification and distribution of watermelon mosaic virus 1 in El Salvador, 437; wilt resistant watermelon cultivars show differential resistance to Fusarium oxysporum f. sp. niveum, 1022
Cupressus arizonica (Arizona cypress): control of Cercospora sequoiae in nursery plantings, 558
---: macrocarpa (Monterey cypress): a preliminary study of the relationship between Coryneum cardinale and Laspeyresia cupressana, 992
Curvularia spp.: 59
--- trifolii f. sp. gladioli: 995
--- intermedius: 192
Cutting decay: of ornamental foliage plant, Scindapsus aureus, found to be major limiting factor in plant propagation in Florida, 1074
Cyclamen: cyclamen wilt, caused by Fusarium oxysporum f. cyclaminis, apparently the first report of occurrence in the Western Hemisphere, 493
Cyperus spp.: common rice field weeds that must be controlled where tests for nematode control effects on rice yields are conducted, 420
Cypress: Arizona cypress, seedling tip blight infection of nursery seedlings, 1192; Monterey cypress, relationship between Coryneum canker and Laspeyresia moth, 992
Cytoplasm: blight ratings and stalk lodging in corn trials in Missouri were greater for

T than for N cytoplasm hybrids, 11; of 25 sources of cytoplasmic male sterility in corn only Tcms or genetically restored TcmsRf lines gave positive reaction to Helminthosporium maydis toxin by leaf whorl bioassay, 29; corn inbred lines with N cytoplasm with different genotypes had varying levels of resistance to H. maydis, 112; reactions of corn seedlings with different cytoplasms to H. maydis toxin, 432; corn in several cytoplasms rated for tolerance to SCLB in the 1971 interregional and regional leaf blight rating tests, 485; results of pairing corn double crosses differing only in type of cytoplasm indicates that both the male and female cytoplasm affected the reaction to seedling diseases, and should be considered in breeding for resistance, 575; sorghum does not show cytoplasmic vulnerability to race T or O isolates of H. maydis, 682; normal, population survival studies of race T and O of H. maydis on a corn hybrid in normal cytoplasm demonstrate the superior ability of race O to become the dominant species, 889; T cytoplasm sweet corn has been found infected with H. maydis in Florida, 969

Cytospora sp.: 108

Damping-off: of tomato transplants was controlled by dipping 'Walter' plants in Difolatan or zinc + maneb suspensions, 840

Dandelion (Taraxacum officinale): yellows disease found in Arkansas, 270

Dasheen (see Colocasia esculenta)

Decay rate: in oaks following fire wounds in North Carolina and Maine, 406

Defoliation: defoliated oak and maple suffered dieback of regrowth by anthracnose, Gnomonia and Gloeosporium, 1046

Delaware: 736

Dendrophoma obscurans: 23

Dermea pseudotsugae: 843

Dew period: dew period, temperature, relative humidity and air flow in relation to spread and increase of southern corn leaf blight, 104; its relation to infection of a male-sterile corn hybrid with race T of Helminthosporium maydis, 767

Diaporthe citri: 80

Dichondra repens: apparently the first report of a mosaic disease of dichondra caused by cucumber mosaic virus, 27

Dieback: of Russian olive, caused by a species of Phomopsis, 995

Dionaea muscipula (Venus-Fly-Trap): foliar blight caused by Colletotrichum gloeosporioides, 391

Dioscorea floribunda: control of tuber rot with benomyl, in India, 990

Diplodia natalensis: 323; 751

Dips: Chemical dips of fungicides ineffective in control of postharvest decay of blueberries, 118; heated fungicidal dips of benomyl or thiabendazole controlled sweetpotato scurf, 122; bare-root dips on Chinese gooseberry cuttings controlled root-knot nematodes, 850

Disease monitoring: for southern corn leaf blight, in 1971, 485

Diseases: fungal and bacterial diseases in the Khuzestan Province of southwestern Iran, 1067

Distribution: of virulence of wheat leaf rust in North Dakota in 1971, 77; of three species of dwarf mistletoe on their principal pine hosts in the Colorado Front Range, 223; peanut rust in Georgia in 1971 was the most widespread on record, 290; of Pythium aphanidermatum in Arizona, 507; of elm phloem necrosis in Pennsylvania, 525; of Dutch elm disease in Colorado in 1971, 912

Ditylenchus dipsaci: 329

Dogwood (Cornus florida): natually infected with tobacco ringspot virus in Maryland, apparently the first report of a virus from dogwood, 336

Douglas-fir (see Pseudotsuga)

Downy mildew: of lima bean, multiple resistance in lima bean breeding lines, 65; of lima bean, strain C of Phytophthora phaseoli found twice in New Jersey in 1971, 237; of sunflower, inbred line HA 61 is valuable source of downy mildew resistance, 428; on St. Augustinegrass in Arkansas, 658; on sorghum, in India, 740; of small grains and sorghum in Arizona, 773

Drop: of lettuce, caused by Sclerotinia sclerotiorum, possible sources of resistance found in some accession lines field-tested in Wisconsin, 875

Drought: as a factor in low level of southern corn leaf blight development in Iowa in 1971, 866

Dutch elm disease (DED): municipal control of DED in Illinois cities, 460; control with soil-amended fungicides, 604; progression of internal and external symptoms, 667; foliar applications of benomyl plus an adjuvant to mature American elms reduced incidence of DED, 685; investigation of elm clons based on physiological vigor for crown resistance to DED, 725; distribution in Colorado, in 1971, 912; continues to move east in the Idaho-Oregon area, 939

Eastern redcedar (see Juniperus virginiana or Cedar)

East Europe: 462
Echinochloa colona: weed involved in nematicide-weed interaction in rice fields, 420
Egypt: 878
Electron microscopy: cadang-cadang-diseased coconut palm leaves revealed viruslike particles in some chloroplasts, 643
Elm: American: elm seedlings grown in sand or perlite planting media took up thiabendazole in higher concentration than those grown in peat, 855
---: Dutch elm disease (DED): 460; 604; 667; 685; 725; 913; 959
---: phloem necrosis: in slippery elm and American elm in 18 counties of New York, 159; distribution in Pennsylvania, 525
---: Siberian (Ulmus pumila), Siberian elm canker observed in all South Dakota counties surveyed, and probably occurs in all counties of the State, 572
El Salvador: 437
Endocronartium harknessii: 1058
Endogone sp.: 58
Endothia gyrosa: 108; 300
England: 770
Epidemic: effect of inoculum density on the development of natural rust epidemics in two wheat nurseries in Minnesota, 410
Epidemiology: of monogenic fungicide tolerance with the plant pathogen Cochliobolus carbonum, under field conditions, 164
Epiphytology: of stripe rust of wheat in northeast Oregon during 1971, 1071
Epiphytotic: of ryegrass blast in Louisiana, 157; of Pyricularia infection on annual ryegrass in Mississippi, 210; of elm phloem necrosis in Finger Lakes, 159
Ergot: of sorghum and millet in India, 740
Erwinia amylovora: 108; 844
--- carotovora: 552
--- --- var. chrysanthemi: 480
--- nimipressuralis: 108
--- tracheiphila: 949
Erysiphe spp.: 852
--- cichoracearum: 45; 878
--- graminis: 75; 358; 618
--- trina: 313
Extraction: of Tylenchulus semipenetrans from citrus roots, by the plastic bag incubation technique, 186

Fern: the tobacco mosaic replication on fern species indicates a wide scope of relationship between TMV and the plant kingdom, 1010
Ferneries: natural oak sites as overstory for ferneries show decline due to cultivation practices that wound or prune the oak roots, 811
Fertilizer: a complete fertilizer, and fertilizer + a preplant nematicide treatment increased yields of sour cherry trees over untreated, 141
Ficus elastica (rubber plant): inoculation of young Ficus leaves with Aphelenchoides besseyi was more reliable, and easier than with older leaves, 471; artificial and natural infection with Cercospora disease, in Florida, 1091
Field evaluation: of triarimol and sulfur fungicides for powdery mildew control on cucumber, 206; of several fungicides for control of Scindapsus aureus cutting decay caused by Pythium splendens, 1074
Field resistance: of 53 Kentucky bluegrasses, mostly from Alaska to powdery mildew in New York, 618
Fire damage: rate of decay in oaks following fire, in North Carolina and Maine, 406; Rhizina root rot of Douglas-fir seedlings planted on burned sites in Washington, 407
Flax: flax rust effectively controlled with oxycarboxin and U8342, 737
Florida: 23; 45; 85; 135; 211; 243; 276; 362; 563; 586; 656; 671; 695; 759; 811; 822; 885; 894; 909; 939; 967; 969; 998; 1019; 1074
Fluorescence: cotton fiber samples examined from 54 locations in U.S. to detect fluorescence associated with Aspergillus flavus boll rot, 215
Fomes annosus: 66; 108; 498; 958; 987
Forage species: variation in disease incidence of five forage species planted on north- and south-facing slopes, 371
Forest tree nurseries: systemic fungicides tested were no better than the formulation routinely used for control of fusiform rust of loblolly pine, 628
Forsythia (Forsythia ovata): natural infection by tobacco ringspot virus, in Maryland, 336
France: 293
Fuirena sp.: common weed in rice fields that is host of ring nematode, 420
Fungi: storage, the lower limit of moisture that permitted invasion of stored sunflowers by Aspergillus glaucus or A. restrictus was about 6.0%, 173
Fungicides (see also Adjuvants, Antibiotics, Chemicals, and Nematicides): effectiveness of four seed-treatment fungicides for rice were significantly improved by treatment of fungicide-treated seed with candidate insecticide PP-211, 154; control of fungicide tolerance to cadmium succinate and cycloheximide by single genes in isolates of Cochliobolus carbonum was not evident under field conditions, 164
---: several tested together: 41 fungicides

evaluated for chemical control of black root rot of bean and tobacco, 15; fungicides as seed treatments reduced the incidence of seedborne Helminthosporium stripe of barley, 183; systemic fungicides triarimol, benomyl and TD-1771 (thiophanate-methyl) applied during growing season controlled stripe smut on Kentucky bluegrass 'Merion', 334; basic copper sulfate was the best of several fungicides tested for control of Arizona cypress blight, with the curtailment of mercury fungicides, 558; ammonium sulfate + chlordane gave best control of Ophiobolus patch disease on 'Highland' bentgrass turf in western Washington, 565; for control of four species of Helminthosporium, 612; for control of flag smut of wheat, 636; benomyl, Difolatan, oil, TBCS and zineb controlled foliar as well as fruit rind infection by greasy spot fungus of citrus, 671; for control of Phytophthora blight of macadamia racemes, 689; for control of powdery mildew of cantaloup, 935; for control of stripe smut of Kentucky bluegrass turf, triarimol was most effective, 1083

---: BAS 3191 F, in single application eliminated Ustilago striiformis and Urocystis agropyri in diseased 'Merion' Kentucky bluegrass, 252

---: BAS 3201 F, controlled powdery mildew on strawberry leaves, 235

---: BAY 79770, controlled powdery mildew of wheat, 358

---: benomyl: young carnation plants treated with benomyl soil amendments had more Alternaria leaf spot than untreated plants, 9; controlled black root rot of bean, and tobacco, 15; controlled Cercosporella foot rot of wheat in Washington, 20; applied weekly or every 2 weeks gave excellent control of strawberry leaf blight, 23; effective control of powdery mildew of orchardgrass, 75; benomyl, captan, and DCNA fungicides were ineffective when used as chemical dips or as preharvest fungicides to reduce postharvest decay of blueberries, 118; controlled sweetpotato scurf, 122; Cercosporella foot rot of winter wheat was controlled effectively at three seeding dates, 178; control of powdery mildew of strawberries, 235; 368; for control of stem blight of Vinca minor, 238; as a substitute for thiabendazole to prevent citrus fruit decay by thiabendazole-resistant strains does not give satisfactory control of resistant strains of blue mold or super-resistant strains of green mold, 256; as a soil drench in glasshouse experiments suppressed invasion of barley roots by Heterodera avenae and delayed subsequent larval development, 261; benomyl and TPTH reduced severity of stem-end blight of pecan, 276; 1st extensive use on peanuts in Georgia may have contributed to increase in rust infection in 1971, 290; preharvest fungicide treatments of strawberries with benomyl and thiophanate-methyl resulted in yield increases, 296; control of foot rot of winter wheat was dependent on rate and time of application, 342; benomyl, chlorothalonil and zinc controlled foliar blight of Dionaea muscipula, 391; benomyl and folpet controlled powdery mildew of kalanchoe, 405; benomyl, chlorothalonil, and thiabendazole are highly effective for control of Rhizoctonia blight of Carissa grandiflora, 511; sprayed on summer plant beds controlled anthracnose of strawberry, 515; reduced number of stunt nematodes parasitizing creeping bentgrass and improved control action of two nonfumigant nematicides, 546; as seed treatment, controlled flag smut of wheat, 636; foliar spray on American elm reduced Dutch elm disease incidence, 685; benomyl and thiophanate-methyl gave control of black root rot of tobacco in New Zealand seedbeds, 708; benomyl and thiabendazole were ineffective for control of Alternaria citri on citrus fruit, 909; effectively controlled pecan scab when applied as spray at 2-, 4-, and 6-week intervals, 980; benomyl applied to cut tuber pieces of Dioscorea floribunda, followed by dusting cut ends with benomyl, controlled tuber rot, 990; was superior to thiophanate-methyl. PMA and dodine in Chile in control of apple scab, 1079; four sprays of 0.10% and 0.05% gave best control of tikka (Cercospora) leaf spot of peanut in India, 1099

---: Bordeaux, two summer applications controlled brown spot needle blight on Scotch pine Christmas trees, 99

---: captan: controlled Thielaviopsis basicola on bean and tobacco, 15; ineffective as chemical dip to reduce postharvest decay of blueberries, 118; captan and DCNA combination as seed treatment reduced Helminthosporium stripe of barley, 183

---: carboxin: carboxin and thiram combination used as seed treatment reduced incidence of Helminthosporium stripe of barley, 183; control of fusiform rust of loblolly pine, 628; effective as seed treatment against flag smut of wheat, 636

---: CELA W 524: orchard control of diseases and mites in apple with new piperazine derivative fungicide in New York, 451;

control of brown rot of plum in New York, 456
---: chloroneb, not effective as seed treatment for Pythium blight of annual ryegrass, 1043;
---: chlorothalonil: controlled Sirococcus tip blight on Jeffrey pine seedlings, 241; for control of foliar blight of Dionaea muscipula, 391; controlled Rhizoctonia blight of Carissa grandiflora, 511
---: DBCP, effective in control of stunt nematode on azalea, 540
---: DCNA: ineffective as chemical dip or preharvest fungicide for postharvest decay of blueberries, 118; DCNA and captan combination reduced Helminthosporium stripe of barley when used as seed treatment, 183; gave effective control of wet-rot of pole bean, 967
---: DD-MENCS, paper or polyethylene sheets as mulch increased efficacy of DD-MENCS in control of Fusarium wilt of tomato, 953
---: Dexon, to control Scindapsus aureus cutting decay, in Florida, 1074
---: Difolatan: Difolatan and zinc + maneb controlled tip blight of Jeffrey pine seedlings, 241; Difolatan was the only one of 10 fungicides tested that provided outstanding control of Phytophthora blight of macadamia, 689; Difolatan or zinc + maneb controlled damping-off of 'Walter' tomato transplants, 840
---: DMTT, controlled black root rot of bean and tobacco, 15
---: EL-279, controlled Urocystis agropyri and Ustilago striiformis on 'Merion' Kentucky bluegrass under greenhouse conditions, 55; reduced ozone injury on bean, 386
---: ETMT: ETMT with thiabendazole were effective against black root rot of bean and tobacco, 15; for control of Pythium blight of ryegrass, 1043; ETMT and thiabendazole were taken up by tomato seedling roots and moved unaltered into stems and leaves, 1054; ETMT was initially superior but Dexon had more lasting effect in control of Scindapsus aureus cutting decay in field tests in Florida, 1074
---: ferbam: ferbam and triarimol not effective as dips for control of sweetpotato scurf, 122; ferbam, carboxin and oxycarboxin gave better control of fusiform rust of loblolly pine than benomyl or chloroneb, 628
---: folpet: controlled powdery mildew of kalanchoe, 405; is ineffective as a fungicide to control rose rust, 1069
---: maneb: controlled Thielaviopsis basicola on bean and tobacco, 15; controlled Lophodermium needle cast of Scotch pine in North Carolina State Forest Nursery; 457; maneb and sulfur/maneb powder formulations reduced pycnidial coverage caused by Septoria tritici on wheat leaves and increased yield, in Israel, 983
---: methyl bromide, methyl bromide and chloropicrin evaluated for nematode and fungi control in celery seedbeds, 661
---: 2-methyl-5,6-dihydro-4H-pyrane-3-carbonic acid anilide (HOE 2989), as seed treatment controlled flag smut of wheat, 636
---: Morestan: Morestan and BAY 79770 controlled powdery mildew of wheat, 358; provided effective control of powdery mildew of cantaloup with high acaricide properties, 935
---: oxamyl (see DPX 1410 under Nematicides)
---: oxycarboxin: almond, chestnut and peach seedlings treated with carbon-labelled oxycarboxin translocated the fungicide more readily when seedlings had been pretreated with two growth retardants, SADH and CCC, 590; control of fusiform rust on loblolly pine, 628; controlled flag smut of wheat when applied as seed treatment, 636; oxycarboxin and U8342 effective in control of flax rust, 737
---: PCNB with granular formulations of V-C 9-104, BAY 68138, MBR 6168, BAY 25141, and D-1410 for control of clubroot and sugarbeet nematode on Brussels sprouts, 375
---: phenylmercuritriethanolammonium lactate reduced occurrence of twig blight stage of sycamore anthracnose in Illinois, 354
---: potassium azide, applied to peanuts after emergence reduced loss of plants from Sclerotium rolfsii but not effective against nematodes, 362
---: PP 675 (Milcurb): gave effective and consistent systemic protection against powdery mildew of squash and cucumber under field conditions in Egypt, 878; as foliar spray gave excellent control of cantaloup powdery mildew, 935
---: SMDC (sodium N-methyldithiocarbamate), controlled black root rot of bean and tobacco, 15
---: sulfur/maneb formulation, reduced pycnidial coverage caused by Septoria tritici on wheat leaves, 983
---: TD-5056 (MSNB) (2-methylsulfonyl-6-nitrobenzothiazole), controlled black root rot of bean and tobacco, 15
---: thiabendazole: controlled Thielaviopsis basicola on bean and tobacco, 15; applied twice weekly, or weekly, controlled Dendrophoma obscurans on strawberry, 23; for sweetpotato scurf, 122; thiabendazole and ETMT were effective in control

of Pythium root rot of Philodendron panduraeforme and Scindapsus aureus, caused by Pythium splendens, 211; shows promise as a "seed" treatment for sweetpotato roots before bedding, 219; controlled Phoma stem blight on Vinca minor, 238; strains of Penicillium italicum and P. digitatum show differential sensitivity to thiabendazole, 256; control of Rhizoctonia blight of Carissa grandiflora, 511; incorporated into soil reduced symptom expression of Dutch elm disease in inoculated trees more than benomyl, captan or Difolatan, 604; uptake in elm seedlings best in sand or perlite medium, reduced in peat, 855; ineffective for control of Alternaria citri on citrus fruit, 909; thiabendazole and ETMT were taken up by tomato seedlings roots and moved unaltered into stems and leaves, 1054
---: thiophanate-methyl (TD-1771): controlled powdery mildew on leaves of strawberries, 235; gave excellent control of stripe smut on 'Merion' Kentucky bluegrass, 334; control of black rot of tobacco, in New Zealand, 708; gave only partial inhibition of apple scab fungus in Chile, 1079
--: thiram: thiram-carboxin combination reduced incidence of Helminthosporium stripe of barley, 183; reduced pycnidial production of Physalospora obtusa on apple mummies, 1102
--: TPTH: reduced severity of stem-end blight of pecan, 276; simultaneous application of TPTH and a sucker control agent to tobacco caused leaf injury, 338
--: triarimol: control of powdery mildew of squash, 45; not effective as dip for sweetpotato scurf, 122; proved superior to sulfur in control of powdery mildew on cucumber, 206; triarimol and EL-279 reduced ozone injury on bean, 386; gave excellent control of powdery mildew of cantaloup, 935; was best of six fungicides for control of stripe smut of Kentucky bluegrass turf in Ohio, 1083
--: triphenyltin chloride, was absorbed and translocated in sugarbeet seedlings in bioassay studies, 776
--: U8342: effective in control of flax rust, 737
--: zinc + maneb: zinc + maneb and benomyl controlled leaf diseases of alfalfa and resulted in increased hay and seed yields, 286; zinc + maneb and cupric hydroxide gave excellent control of Xanthomonas dieffenbachiae on Philodendron, 394; zinc + maneb or Difolatan controlled dampingoff of 'Walter' tomato transplants, 840; was the most effective and cheapest fungicide of four tested in Uganda for control of bean diseases, 901; zinc + maneb-Karathane and thiram reduced pycnidial and conidial production of Physalospora obtusa on apple mummies in Connecticut, 1102
---: zineb: controlled geranium rust in Pennsylvania greenhouse, 759; gave some control of leaf blight of tomato caused by Phoma destructiva, 869; Cercospora disease of Ficus elastica was controlled with zineb, 1091
Fusarium sp.: 108; 318; 575; 661; 942; 1040
--- spp.: 80; 300; 783; 861
--- equiseti: 963
--- moniliforme: 836; 965
--- --- 'Subglutinans': 836
--- nivale: 32; 803
--- oxysporum: 108; 963
--- --- f. sp. cubense: 1000
--- --- f. cyclaminis: 493
--- --- lycopersici: 695; 953; 1054
--- --- f. sp. niveum: 1022
--- --- f. sp. vasinfectum: 328
--- roseum: 42; 788
--- semitectum: 693
--- solani: 310; 523; 600; 944; 963
--- solani f. sp. pisi: 398
Fusicladium effusum: 980

Galling: caused by root-knot nematodes on five varieties of soybean in Egypt, 882
Ganoderma curtisii: 811
Genotypes: corn inbreds showed varying levels of susceptibility and resistance to H. maydis race T, 112
Geographic distribution: of Aspergillus flavus boll rot in experimental plots across the U.S. Cotton Belt in 1970, 215
Georgia: 146; 147; 290; 534; 796
Geotrichum candidum: 582
Geranium: in controlled environments, lesion size and sporulation of Botrytis on intact geranium leaves increased with temperature to 25°C, light appeared to have no significant effect, 126; geranium rust discovered in Pennsylvania, 759
Ghana: 1011
Gibberella zeae: 836
Gladiolus: incidence of the Curvularia disease in Illinois, 995
Gloeosporium aridum: 108
Glomerella cingulata: 108
Glume blotch: of spring wheat, caused by Septoria nodorum, none of 30 spring wheats tested was highly resistant to Septoria, 761
Gnomonia spp.: 1046
--- fructicola: 998
--- nerviseda: 276
--- platani: 354
Gossypium barbadense (see Cotton)

Graminella nigrifrons: 652
Grape: tomato ringspot virus found in 'Baco Noir' grapevines in New York, 133; its rapid spread since 1971, 1062; Hemicriconemoides chitwoodi on grapevines in California, 1028; control of nematode-fanleaf infected vineyards by soil fumigation with 1,3-D and methyl bromide, 1031
Grasses: greenhouse inoculation tests indicate five species of Curvularia are pathogenic to cultivars of Poa pratensis, Festuca rubra, and Agrostis palustris, 59
---: bentgrass: Tylenchorhynchus dubius controlled on 'Toronto' creeping bentgrass with benomyl and nonfumigant nematicides, 546; ammonium sulfate plus chlordane gave best control of Ophiobolus patch disease on 'Highland' bentgrass, 565
---: bluegrass: 56 cultivars of Kentucky bluegrass evaluated for resistance and susceptibility to Typhula blight and Fusarium patch, 'Adorno' was superior to the others, in Michigan, 32; fungicide EL-279 gave excellent control of Urocystis agropyri and Ustilago striiformis in minute dosage on 'Merion', 55; stripe and flag smuts controlled on 'Merion' with BAS 3191F in a single application to soil, 252; triarimol, benomyl and thiophanate-methyl gave excellent control of stripe smut of 'Merion' bluegrass turf when drenched into the root zone, 334; disease incidence as affected by north- and south-facing slopes in West Virginia, 371; five genera of nematodes controlled in bluegrass lawn in Maryland with nematicide V-C 9-104, 476; 55 Kentucky bluegrass introductions, mostly from Alaska, and six named cultivars were tested for field resistance to powdery mildew in New York, 618; Tylenchorhynchus nudus and other nematodes associated with Kentucky bluegrass turf in South Dakota, 898; triarimol was most effective of six fungicides for control of stripe smut on Kentucky bluegrass, 1083
---: bermudagrass: a spotting and discoloration condition of dormant bermudagrass by a sterile basidiomycete, 355; incidence of disease on north- and south-facing slopes in West Virginia, 371
---: centipedegrass (Eremochloa ophiuroides) Fusarium blight caused by Fusarium roseum identified in Alabama, 42
---: fescue: symptom expression of barley yellow dwarf virus in tall fescue, 167; effect of slope orientation on incidence of disease in tall fescue in West Virginia, 371; two races of Puccinia coronata var. coronata were found on tall fescue in Alabama, 614
---: johnsongrass, johnsongrass occurring naturally in Louisiana corn and sugarcane producing areas was surveyed for presence of maize dwarf mosaic virus, 622
---: orchardgrass, benomyl, in greenhouse tests, provided longterm control of powdery mildew, 75
---: ryegrass: Pyricularia sp. caused an epiphytotic of leafspot and blast disease in Louisiana in fall of 1971, a first report of this disease in the State, 157; an epiphytotic caused by a species of Pyricularia occurred in Mississippi in fall of 1971, 210; ETMT applied to seed as slurry or as a dust was effective in control of Pythium blight, 1043
---: St. Augustinegrass: St. Augustine Decline virus found in Louisiana, 69; rust, caused by Puccinia stenotaphri was found on St. Augustinegrass in the Gulf Coast areas of Texas, 630; downy mildew found in trace amounts in Arkansas, 658; in south Florida all nine virus isolates collected from St. Augustinegrass were strain E of sugarcane mosaic virus, 822; characterization of strain E of sugarcane mosaic virus isolated from this grass in Florida, 885
---: zoysiagrass: rust, conditions that aid its development and perpetuation on zoysiagrass, 227; V-C 9-104 nematicide controlled five genera of nematodes in zoysiagrass lawn in Maryland, 476; nematodes contribute to zoysiagrass decline in Maryland, 568
Gray mold: effect of temperature and light on colonization and sporulation of Botrytis cinerea on geranium, 126; of poinsettia, effects of Botrytis and ozone on bracts and flowers, 814
Greasy spot rind blotch: of citrus, caused by Mycosphaerella citri, 671
Greece: 264
Green mold (Penicillium digitatum): differential sensitivity to thiabendazole in spore populations from California lemon packinghouses, 256
Groundnut (see Peanut)
Guatemala: 753
Guava (Psidium guajava): dry rot of guava fruits caused by Diplodia natalensis, 323
Guyana, West Indies: 1059
Gymnosporangium clavipes: 451
--- juniperi-virginianae: 451

Hackberry (Celtis spp.): found to be host of pecan pathogen, Cristulariella pyramidalis, 176
Hawaii: 73; 331; 446; 689; 904
Helicotylenchus spp.: 551; 898; 1059; 1077
--- pseudorobustus: 476

Helminthosporium spp.: as laboratory contaminants on incubated leaf specimens, 834
--- avenae: 648
--- dictyoides: 371
--- gramineum: 183; 891
--- maydis: 11; 29; 104; 112; 203; 247; 432; 612; 659; 660; 682; 767; 805; 889; 969; 1087
--- rostratum: first report of this pathogen on rice in the United States, 26
--- sorokinianum: 788
--- triseptatum: 612
--- turcicum: 11; 612
--- vagans: 371; 612
Hemicriconemoides chitwoodi: 1028
Hendersonula toruloidea: 50
Heterodera avenae: 261
--- schachtii: 375; 432
--- tabacum: 255
Hoplolaimus galeatus: 476; 691
Hormiscium sp.: 275
Host range: of strain E of sugarcane mosaic virus is limited to the Gramineae, 885
Hosts: of Cristulariella pyramidalis, cause of zonate leafspot of pecan, 176
Hosts and diseases: of plants found in Southwestern Iran on the Dez Irrigation Project, 1067
Host-specificity: of a toxin from Helminthosporium maydis race T, 247
Hot water treatments (see Temperature)
Hydroponic tomatoes: bacterial canker and root knot diseases controlled by flushing gravel beds with 2% formalin, 139
Hypoxylon atropunctatum: 300
--- mediterranium: 300

Idaho: 400; 498; 939
Illinois: 354; 460; 995
India: 64; 115; 323; 424; 601; 693; 740; 990; 1099
Indiana: 358; 827
Inoculation: inoculation method using perlite infested with Rhizoctonia solani was the most effective of three methods tested for measurement of R. solani resistance in potatoes, 348; young vs. older leaves of rubber plant for inoculation with foliar nematode, Aphelenchoides besseyi, 471; root chip inoculation for indexing for stem pitting of Prunus, 785; a multiculture inoculation system for study of host: parasite relationships of cereal crop species and rust fungi, 847; inoculation of large numbers of pine and oak hosts of Cronartium fusiforme with specified spore densities, 931; of loblolly stumps with Peniophora gigantea in chain SAE 10 saw oil, 958; of severed roots of eastern redcedar with Fomes annosus, 987
inoculation, incubation and abscission of Ficus elastica foliage with Cercospora sp., 1091
Inoculum density: of stem and leaf rust of wheat and the effect on natural rust epidemics in wheat nurseries, 410
Insecticides (see Chemicals)
Internal disorder: of sweetpotatoes found only in moist-type, consists of hard inedible tissues when cooked, 88
Iowa: 659; 660; 866
Iran: 206; 1067
Iraq: 824
Isolates: variation among Washington isolates of beet western yellows virus, 198; isolate of cocoa swollen shoot virus from Nigeria is different from other isolates studied previously, 755
Israel: 600; 963; 983

Jamaica: 382
Japan: 490
Juniperus virginiana (eastern redcedar): a direct relationship between tree vigor and susceptibility to Fomes annosus in eastern redcedar was seen in severed roots of dominant and suppressed trees, 987

Kalanchoe: folpet and benomyl were effective fungicides for control of powdery mildew, Karathane not effective, 405
Kallstroemia maxima: an annual plant that is little affected by herbicide barriers, maintained to contain burrowing nematode within infested areas in Florida, is a host of this nematode, 85
Kansas: 957
Kenaf (Hibiscus cannabinus): zonate leaf spot, caused by Cristulariella pyramidalis, defoliated all kenaf varieties on trial in southern Mississippi, 917; rust (Aecidium garkeanum) attacked kenaf in pilot plantings and variety trials in Zambia, 1047; susceptibility of kenaf to a new race of anthracnose in Zambia, 1049
Kentucky: 652; 1026
Kiwi fruit (see Chinese gooseberry)
Koa (Acacia koa): collar rot is caused by Calonectria crotalariae, 73

Landscape trees: diseases of shade and ornamental trees in Virginia, with 34 new reports, 108
Laspeyresia cupressana: 992
Lateral root production: of soybean plants as affected by infection with Meloidogyne javanica and M. incognita, 882
Late-spring leaf scorch: of maple and beech trees in southern Ontario, 526
Leaf blight (see Blight)
Leaf blotch: of wheat, caused by Septoria

tritici in Israel, the disease potential is great for substantial yield loss, 983; of strawberry, caused by Gnomonia fructicola, in Florida, 998
Leaf freckles and wilt: of corn, a new corn disease in Nebraska, caused by a previously undescribed species of Corynebacterium, 863
Leaf pigments: in sweetpotato, and their relation to foliar symptom expression of viral infection, 501
Leaf scald: of rice, in West Africa, 106
Leaf spot: of carnation, caused by Alternaria dianthi was more severe on benomyl-treated plants than untreated, 9; Cristulariella pyramidalis caused severe defoliation to silver maple in Florida, 135; zonate leafspot of pecan, hosts found in pecan orchard hedgerows, 176; of alfalfa, Pseudopeziza medicaginis controlled with zinc + maneb and benomyl, 286; of cotton, estimated yield loss in 1971 from Alternaria, Cercospora, Ramularia and other species, 328; of strawberry, relation of temperature to leaf spot caused by Mycosphaerella fragariae, 345; of oat, caused by Helminthosporium avenae, oat varieties and Avena species were compared for reaction to virulent strain of pathogen from Puerto Rico, 648; of cowcockle (Saponaria vaccaria) caused by Alternaria dianthi, in Montana, 728
---: of apple: an Alternaria was found associated with leaf spot of apple, in New York, 721; of 'Cortland' apples, fungicide sprays greatly reduce infection of immature apples, reduce the production of conidia on mummied fruit the following season, and subsequent incidence of frogeye leaf spot, 1102
---: of peanut: 12 of 58 varieties tested under artificial conditions in India were resistant, 23 were tolerant and 23 were susceptible to Cercospora personata and C. arachidicola, 601; caused by C. personata was contributing factor in early harvesting of some fields in Texas in 1971, 873; zonate leaf spot of peanut, caused by Cristulariella pyramidalis, 796
Leaf wetness: its effect on infection of skeleton weed by Puccinia chondrillina, a potential biological control agent, 858
Leptosphaerulina briosiana: 286
Lesion development: by Pyricularia oryzae, resistance of the host cells of rice cultivar Peta is triggered by the invading blast pathogen itself, 960
Lethal yellowing disease: of coconut palms, tetracycline antibiotics cause cessation of symptom development, 1019
Lettuce: ascospore infection by Sclerotinia sclerotiorum is a more important factor in lettuce drop in Wisconsin than was previously realized, 798; five of 170 accession lines of lettuce showed most resistance to lettuce drop in field evaluation in Wisconsin, 875; some factors that influence the susceptibility of lettuce to CO_2 injury, 1051
Light: light had no apparent effect on colonization or sporulation of Botrytis on geranium leaves in light vs. darkness at 20°C, 126
Light intensity: light shade or direct sunlight give better leaf symptom expression in virus-infected sweetpotatoes, 501
Lilac (Syringa oblata affinis): lilac mottle suggested as name for virus isolated from lilac in Maryland, 923
Limonium perezii (see Statice)
Lissorhoptrus oryzophilus: 154
Lodging: of corn, stalk lodging increased as leaf blight rating increased in Missouri trials, 11
Longevity: of Phytophthora parasitica chlamydospores in water, 971
Longidorus siddiqii: 699
Longterm control: of powdery mildew of orchardgrass with benomyl, 75
Lophodermium pinastri: 457
Losses: survey to assess economic losses of food, fiber, and ornamental crops in Pennsylvania from air pollution, 701
Louisiana: 58; 69; 157; 272; 276; 368; 420; 515; 622

Macadamia: field bioassay and evaluation of fungicides for control of Phytophthora blight, 689
Macrophomina phaseoli: 115
Madrone (Arbutus menziesii): unreported canker disease of Madrone, caused by Hendersonula toruloidea found in Washington State, 50
Maine: 406
Maize (see Corn)
Male-sterile cytoplasm (see Cytoplasm)
Mango: postharvest disorders of mangos as affected by fungicides and heat treatments, 751
Maple: late-spring leaf scorch is caused by sudden exposure to full sunlight after wet cloudy period, 526
---: red, regrowth of manually defoliated red maple trees became infected by Gloeosporium sp. in Connecticut, 1046
---: sugar: Fusarium canker, levels of precipitation were not correlated with canker frequencey, 942; study on maple stem tissue extracts indicates there are factors other than the presence of water-soluble compounds that either inhibit or stimulate

spore germination and apparently determine the resistance of sugar maples to Fusarium solani, 944; effect of salt injury on shoot growth of sugar maple, 1004
Marketability: of summer squash, watermelon mosaic causes loss of marketable fruits, 147
Market diseases: Aspergillus rot of limes in local markets in India, 64; internal disorder of sweetpotatoes sampled in Greater New York market, 88
Maryland: 336; 476; 568; 923
Massachusetts: 832
Mealypod disease: of Theobroma cacao, was observed in Ghana cocoa-growing areas between June and November 1971, but not from December 1971 to April 1972, 1011
Melampsora abietis-canadensis: 108
--- lini: 737
Melanconis juglandis: 108
Meloidogyne sp.: 328; 551; 1059; 1077
--- spp.: 169
--- arenaria: 824
--- hapla: 417; 520; 850
--- incognita: 139; 618; 824; 882; 914
--- --- acrita: 305; 1065
--- javanica: 882; 914
Methods and techniques: a method of collection of ascospores of Venturia inaequalis on glass slides at selected sites and in controlled numbers for quantitative studies in spore germination, 39; isolation of Phytophthora palmivora from bark cankers of cocoa without using cocoa pods, 71; root incubation technique for extraction of citrus nematodes from roots incubated at 70°F in plastic bags containing 10 ml of 3% hydrogen peroxide, 186; a reliable and rapid method for detecting contamination of potato tubers by Erwinia carotovora, 552; paste application of nematicide to citrus seedling stems to control Pratylenchus coffeae, 626; method described to make potential users of rotorod impaction aerosol samplers aware of the importance of considering sampling efficiency, 704; for extraction of nematodes from heavy clay soils, sodium hexametaphosphate is used to disperse the clay particles, followed by direct centrifugation in sugar solution, 808; use of black strap molasses as a sugar substitute for nematode extraction from soil by the flotation-sieving method, 1093
---: equipment, cylindrical rods are more efficient spore samplers than glass slides, 719
---: inoculation: three inoculation methods to screen potatoes for resistance to Rhizoctonia solani, 348; root chip inoculation to index on peach seedlings for stem pitting, 785; a multi-culture inoculation system to test wheat leaves with four uredial cultures of rust at the same time, 847; an improved method for large-scale inoculations of pine and oak with Cronartium fusiforme, 931
Mexico: 803; 896; 1038
Michigan: 32; 275; 378
Microclimate: of northern and southern exposures in relation to development of specific plant pathogens, 371
Microsphaera alni: 313; 852
Millet: India surveyed for diseases of sorghum and millet in 1970, 740
Mississippi: 210; 916; 917
Missouri: 11; 167
Mistletoe: dwarf, distribution of three species of dwarf mistletoe on their pine hosts in the Colorado Front Range, 223
Moisture content: 6% moisture in stored sunflower seeds was the lower limit for prevention of invasion by storage fungi, 173
Molasses: as an economical substitute for sugar in the extraction of nematodes from soil by the flotation-sieving technique, 1093
Momordica charantia: source of watermelon mosaic virus 1 for cucurbit crops in Palm Beach County, Florida, 563
Monilinia fructicola: 456
Monilochaetes infuscans: 122; 219
Monogenic tolerance: of Cochliobolus carbonum isolates to fungicides cadmium succinate and cycloheximide field-tested in Pennsylvania, 164
Montana: 188; 728; 761
Montserrat: 817
Mukwa (Pterocarpus angolensis): mukwa blight is killing these trees of economic importance (for furniture) in Zambia, Central Africa, 820
Mulching: use of paper or plastic mulches without fumigation greatly retarded Fusarium wilt development of tomato in Florida tests, 953
Mummies: of apple, fungicide sprays greatly reduced Physalospora obtusa pycnidial and conidial production on the mummies the following season, 1102
Municipal control: of Dutch elm disease continues to be effective in Illinois cities, 460
Mycoplasma: remission of lethal yellowing of coconut palm with tetracycline antibiotics lends credence to the hypothesis that the disease is of mycoplasmal etiology, 1019
Mycosphaerella citri: 671
--- fragariae: 345
Myzus persicae: 441; 597; 731

Nebraska: 863
Needlecast: of Pinus albicaulis and P. linearis, caused by Bifusella saccata and B. linearis in California's Sierra Nevada, 102; of Scotch pine, caused by Lophodermium pinastri, controlled with maneb, 457
Nematicides: (see also Chemicals and Fungicides): for nematode control in rice fields, nematicides may stimulate grassy weed development, 420
---: BAY 25141 (Dasanit): tested for control of sugarbeet nematodes on Brussels sprouts, 375; benomyl enhanced activity of BAY 25141 for control of stunt nematode on creeping bentgrass, 546; was more effective than DBCP in suppression of nematodes in zoysia turf, 568
---: BAY 68138 (Nemacur): for control of sugarbeet nematode on Brussels sprouts, 375; a 10% paste applied to stems of citrus seedlings infected with Pratylenchus coffeae reduced nematodes in roots, 626
---: cynem, controlled root-knot nematode on Chinese gooseberry cuttings, 850
---: 1,3-D: as preplant fumigant for apple tree orchards in New York, 577; high rates and deep placement of 1,3-D have controlled Xiphinema index and fanleaf virus for 5 years in replanted vineyards, 1031
---: DBCP: fumigated rows (preplant) of peach planting sites with DBCP resulted in less stem pitting than in untreated rows, in New Jersey, 162; ethylene dibromide and DBCP as preplant treatment reduced root-knot index on peach trees in South Carolina, 169; DBCP, 1,3-D, and MBR-CP as preplant fumigants for apple tree orchards in New York, 577; applied with double chisel to narrow double-row cotton in Texas increased yield and nematode control over fumigation with one chisel to single wider rows, 1065
---: DPX 1410 (oxamyl): as foliar spray controlled Heterodera tabacum, 255; showed poor control of sugarbeet nematode but yield of Brussels sprouts increased, 375; in combination with benomyl improved control of stunt nematode on creeping bentgrass, 546
---: ethylene dibromide, reduced root-knot index on peach trees in South Carolina, 169
---: methyl bromide, at 400-800 lb/A under continuous polyethylene sheeting shows promise as a control for Xiphinema index and fanleaf virus, 1031
---: MBR-CP: preplant fumigation for apple orchards in New York resulted in increased tree growth, 577; evaluated for nematode and soil fungi control in celery seedbeds on muck soil, 661
---: Tirpate (MBR 6168), initial trial showed excellent control of sugarbeet nematodes on Brussels sprouts, 375
---: V-C 9-104 (Mocap): controlled sugarbeet nematode on Brussels sprouts, 375; controlled several nematode genera in lawngrass in Maryland, 476; controlled root-knot nematodes on Chinese gooseberry cuttings, 850
Nematodes: Fusarium sp. and a nematode associated in new citrus disease complex in Tanzania, 318; yield loss of cotton caused by nematodes in the U.S. in 1971, 328; red ring disease, of coconut palm, caused by Rhadinaphelenchus cocophilus, in West Indies, 339; direct centrifugal-flotation method for extraction of nematodes from heavy clay soil, 808; Hemicriconemoides chitwoodi tested for host suitability to 15 kinds of plants but reproduction occurred only on Vitis vinifera, V. rupestris and 1613 (Solonis x Othello), 1028; molasses instead of sugar in the preparation of extraction solution for the flotation-sieving process for nematodes, 1093
---: burrowing, an annual herb Kallstroemia maxima found to be new host for Radopholus similis, 85
---: citrus: Tylenchulus semipenetrans, root incubation technique for extraction from citrus roots, 186; evaluation of citrus selections to T. semipenetrans, 747
---: cyst: sugarbeet cyst nematode, Heterodera schachtii, population levels and damage to red table beet and cabbage under greenhouse conditions, 434; controlled on Brussels sprouts with low vapor nematicides, 375
---:---: tobacco cyst nematode, controlled with nematicide-insecticide D-1410, 255
---: dagger, Xiphinema americanum, zoysiagrass decline, 568
---: lance, Hoplolaimus galeatus, zoysiagrass decline, 568; two races of sand pine (Pinus clausa) show no resistance to H. galeatus in inoculation studies, 691
---: lesion: Pratylenchus penetrans, preplant nematicide treatment + plant nutrients increased yields of cherry trees over untreated trees, 141; P. penetrans is pathogenic to many ornamental plants in Arizona, 914
---: needle, pathogenicity of Longidorus siddiqii on Egyptian cotton, 699
---: plant-parasitic: nematodes associated with some economic crops in Guyana, 1059;

preliminary study of nematodes associated with selected root crops at the University of the West Indies, 1077
---: reniform: found associated with unthrifty papaya trees in Trinidad, 551; the role of Rotylenchulus reniformis in the development and severity of Fusarium wilt of cotton, 1040
---: ring: Criconemoides sp. not controlled with potassium azide broadcast postemergence on peanuts, 362; control of C. onoensis in rice fields with nematicides and the interaction of weeds, 420; cause of zoysiagrass decline, 568
---: root-knot: control of root-knot nematodes in hydroponically grown tomatoes by flushing gravel beds with 2% formalin, 139; preplant fumigation resulted in higher yields on peach trees in South Carolina, and low root-knot index, 169; 30°C appeared an optimal temperature for survival of Meloidogyne incognita acrita in roots of 'Nemaguard' peach seedlings, 305; potassium azide did not control root-knot nematodes in peanut fields when broadcast after emergence, 362; survival of root-knot nematodes on roots of rhubarb in a tobacco greenhouse, 417; five biotypes of M. hapla established in Pacific Northwest, 520; on papaya, in Trinidad, 551; M. incognita found pathogenic to Colocasia esculenta in Trinidad, 618; M. incognita and M. arenaria identified in Iraq for first time, 824; M. hapla controlled on Chinese gooseberry with bare-root dip in nematicides, 850; M. incognita and M. javanica found capable of parasitizing many common ornamentals in urban areas of Arizona, 914; double chisel DBCP fumigation of cotton planted in double rows showed yield increase over single chisel fumigation of wider rows, 1065
---: spiral: on papaya, in Trinidad, 551; Helicotylenchus pseudorobustus and zoysiagrass decline, 568
---: stem, onions screened in controlled environment for resistance to Ditylenchus dipsaci, 329
---: sting, Belonolaimus longicaudatus reported from Kansas, 957
---: stubby root, Trichodorus christiei not effectively controlled with DBCP drenches on azalea, 540
---: stunt: control on azalea with DBCP drenches, 540; Tylenchorhynchus claytoni associated with zoysiagrass decline, 568; Tylenchorhynchus nudus was present in 46% of samples from Kentucky bluegrass turf in South Dakota, Paratylenchus occurred in 98% of samples, 898
New Brunswick: 245
New diseases: a mosaic disease of Dichondra repens found caused by cucumber mosaic virus, 27; canker disease of Madrone, caused by Hendersonula toruloidea, in Washington State, 50; Pythium root rot of Philodendron panduraeforme and Scindapsus aureus, 211; bacterial stem rot of poinsettia, caused by Erwinia carotovora var. chrysanthemi, in Ohio, 480
New distribution: 34 new reports of diseases of shade trees in Virginia, 108
---: bacterial blight of carrot in Wisconsin, 744
---: bacterial leaf spot of clover caused by Pseudomonas stizolobii found in Hawaii, 446; bacterial leaf spot of onion caused by P. syringae found in Japan, 490
---: of Bifusella saccata and B. linearis on whitebark pine in mountainous regions of California, 102
---: blast disease of ryegrass found in Louisiana, 157
---: cyclamen wilt in California, first report of occurrence in Western Hemisphere, 493
---: Dermea pseudotsugae found in California on young Douglas-fir, 843
---: Fusarium nivale on triticale in Mexico, at Toluca, 803
---: geranium rust Puccinia perargoniizonalis found in Johnstown, Pennsylvania in 1971, 759
---: Helminthosporium rostratum found on rice in the United States for the first time, Myrtle Beach, South Carolina, 26
---: leaf scald of rice, caused by Rhynchosporium oryzae, identified in Nigeria for first time, 106
---: Meloidogyne arenaria and M. incognita found in Iraq, 824
---: peanut stunt virus reported for first time from Alabama, 415
---: of Phomopsis root rot of cucumber near Vancouver, British Columbia, 53
---: Phytophthora root rot of alfalfa in Washington, 830
---: Rotylenchulus reniformis found in Trinidad, apparently a first report, 551
---: first report of Scleroderris largerbergii in New Brunswick, Canada in 1971, 245
---: St. Augustine Decline virus reported from Louisiana for the first time, 69
---: stem pitting found in sour cherry in Poland, 462
---: target spot of tomato, not previously reported in U.S. literature, discovered in Florida, 243
---: apparently the first report of occurrence

of watermelon mosaic virus 1 in Central America, 437
---: zonate leafspot caused injury to 10 deciduous hosts in Florida, new hosts records on 8 genera, 135; Cristulariella pyramidalis found in Mississippi on kenaf plantings, 917
---: first report of Xanthomonas dieffenbachiae in Hawaii, on anthurium, 904
New Hampshire: 1004
New hosts: bean yellow mosaic virus on Proboscidea jussieui, 548
---: Bifusella saccata on Pinus albicaulis in California, 102
---: Erysiphe trina on Lithocarpus densiflora and Quercus chrysolepis, in California, 313
---: of Coleosporium crowellii in Western States and Mexico, Pinus cembroides, P. edulis, P. culminicola, and P. monophylla, 474
---: Cristulariella pyramidalis on peanuts (Arachis hypogaea) in Georgia, 796
---: of lettuce mosaic virus, Cucurbita pepo, Iberis umbellata, Melilotus italica, Trifolium incarnatum, Tragopogon porrifolius, and some cultivars of Cucurbita maxima and Phaseolus vulgaris, 281
---: Microsphaera alni on Lithocarpus densiflora, L. densiflora var. echinoides, and Quercus chrysolepis, in California, 313
---: Phyllactinia corylea on Quercus kelloggii, in California, 313
---: red pine is new host for brown spot (Scirrhia acicola) in Wisconsin, 712
---: Sclerotium rolfsii on statice in Florida, 656
---: Sphaerotheca lanestris on Quercus cerris, Q. chrysolepis, Q. dunnii, Q. laurifolia, Q. obtusata, Q. palustris, and Q. velutina, in California, 313
---: tobacco ringspot virus on dogwood (Cornus florida) in Maryland, 336
---: of Verticillium wilt, found in Greece, 264
---: zonate leafspot, caused by Cristulariella pyramidalis, found on eight new hosts in Florida, 135
New Jersey: 85; 162; 237
New York: 133; 159; 281; 285; 434; 451; 456; 548; 577; 616; 618; 721; 1062
New Zealand: 708
Nigeria: 106; 755
North Carolina: 92; 118; 406; 457; 774
North Dakota: 77; 428; 788

Oak: four species of powdery mildews occur on oaks in the San Francisco Bay area of California, 313; black and scarlet oaks, difference in decay rate after fire, between oaks in North Carolina and Maine, 406; decline, root pruning and wounding during cultivation of ferneries provides infection courts for root and butt decay fungus, Ganoderma curtisii and the root decay fungus, Clitocybe tabescens, 811; predisposition of oak (Quercus agrifolia) to powdery mildew infection is genetic as well as physiologic, 852; northern red oak (Quercus rubra), a rapid inoculation method for large numbers of pine and oak hosts with known inoculum sources of Cronartium fusiforme at specified spore densities, 931; red oak (Quercus rubra), attack on regrowth of defoliated oak trees by Gnomonia spp., 1046
Oat: tolerance in oat varieties to barley yellow dwarf virus, 230; stem rust, race 31 comprised 63% of 937 isolates collected in the U.S. in 1971, 608; leaf spot (Helminthosporium avenae), Avena spp. tested for reaction to a virulent culture of H. avenae from Puerto Rico, 648
Ohio: 523; 536
Oil: chain oil of power saw, as carrier of Peniophora gigantea to control Fomes annosus, 66; chain oil SAE 10 instead of SAE 30 resulted in excellent oidial germination of Peniophora gigantea, a carrier to control Fomes annosus invasion of pine stumps, 958
Oklahoma: 35; 980; 1022
Olive: Phomopsis dieback of Russian olive, 995
Onion (Allium spp.): resistance to Ditylenchus dipsaci in four onion species, several selections exhibited resistance, 329; bacterial leaf spot, caused by Pseudomonas syringae identified in Japan, 490
Ontario: 417; 526
Ophiobolus graminis: 565
Oregon: 178; 180; 939; 1071
Ornamentals: Phytophthora cinnamomi found in South Australia extensively in Adelaide Botanic Garden properties in ornamental nurseries, mainly from the families Proteaceae, Myrtaceae and Ericaceae, 290; kalanchoe powdery mildew controlled with benomyl and folpet, 405; Carissa grandiflora, a tropical plant developed water-soaked spots on leaves, caused by Rhizoctonia solani, 511; various Arizona-grown ornamentals found susceptible to attack by nematodes, 914; Tecomaria capensis (cape honeysuckle) found naturally infected with alfalfa mosaic virus, in Israel, 1035
Ovulinia azaleae: 1011
Ozone: hydrocarbon wax emulsion sprays at 5 or 10 ml/liter water protected bean plants from ozone injury, 6; ozone had little effect on chrysanthemum varieties, 85;

relative tolerances of 12 safflower cultivars to ozone, 195; suppression of ozone injury to Phaseolus vulgaris with triarimol and its analogue, 386; poinsettia plants exposed to ozone for 4 hr at levels up to 45 pphm had no measurable effect on incidence of Botrytis on flowers and bracts inoculated just prior to ozone exposure, 814; sensitivity of spinach cultivars to ozone, 832

Pacific Northwest: 520; 792
Palm: coconut, a survey of red ring disease in Grenada, West Indies, 339
Panonychus ulmi: 451
Papaya: spread of papaya ringspot virus in Hawaii, 331; Rotylenchulus reniformis found associated with papaya roots in Trinidad, 551
Paratylenchus sp.: 898
Pathogenicity: of Curvularia spp. to turfgrass, 59; of Longidorus siddiqii on Egyptian cotton in pot culture, 699; of Fusaria isolated from avocado, banana and citrus fruit in Israel, 963; watermelon cultivars compared for differential resistance to Fusarium wilt in Oklahoma trials, 1022
Pea (Pisum sativum): three plant introduction lines had resistance to pea fizzletop disease, 131; artificial infestation of large pea field plots with Fusarium solani f. sp. pisi in Washington, 398
Peach: nematodes: peach trees planted on fumigated soil or on resistant rootstocks in Meloidogyne-infested soil in South Carolina produced greater yields and greater growth, 169; influence of temperature on infection of root-knot nematodes on 'Nemaguard' seedlings, 305
---: stem pitting: incidence in fumigated and nonfumigated field plantings in New Jersey, 162; stem pitting found for first time in New York State, 616; root chip inoculation for indexing for stem pitting, 785
Peanut (Arachis hypogaea): post-emergence application of potassium azide to peanuts reduced loss from Sclerotium rolfsii but not effective for nematodes, 362; peanut stunt virus in Alabama, 415; soil management practices for control of Pythium rot of peanut pods in Israel, 600; reaction of peanut varieties against Cercospora personata and C. arachidicola, 601; Cristulariella pyramidalis found on peanut for the first time in 1971 in Georgia, 796
---: rust: distribution and significance of peanut rust in Georgia in 1971, 290; screening for resistance to peanut rust in greenhouse and field, 382; development and spread of peanut rust in south Texas in 1971, 873
Pear: fire blight, incidence of trunk blight in Magness cultivar, 844
Pearl millet (Pennisetum typhoides): Pennisetum strain of maize streak virus found in India on pearl millet, 424
Pecan: trumpet vine, hackberry, Virginia creeper, poison oak, and Sassafras in pecan orchard hedgerows were found to be hosts of Cristulariella pyramidalis, cause of zonate leafspot of pecan, in Alabama and Georgia, 176; fungicides reduce incidence of stem-end blight in Louisiana tests, 276; pecan scab, long-term control with foliar applications of benomyl, in Oklahoma, 980
Penicillium sp.: 990
--- spp.: 861
--- atrovenetum: 1054
--- digitatum: 256
--- italicum: 256
--- implicatum: 919
--- rugulosum: 919
Peniophora gigantea: chain saw-applied oidia of P. gigantea to loblolly stumps to control Fomes annosus colonization, 66; 958
Pennsylvania: 525; 701; 759; 836; 942; 1058
Pepper: naturally occurring pepper virus strains in south Florida, and reactions of pepper cultivars to them, 586; a new virus disease of pepper in Arizona is from the potato virus Y group, 731
Peridermium filamentosum: 896
Persimmon (Diospyros virginiana): outbreak of persimmon wilt caused by Cephalosporium diospyri following damage of Hurricane Camille in Mississippi, 916
Phaeosphaeria herpotricha: 276
Phaseolus lunatus (lima bean) (see Bean)
Phaseolus vulgaris (Pinto III bean) (see also Bean): triarimol and its EL-279 analogue suppressed ozone injury, 386
Philippines: 643; 960
Philodendron oxycardium: streptomycin failed to control Xanthomonas dieffenbachiae on Philodendron apparently because of pathogen resistance to the antibiotic, 394
--- panduraeforme: Pythium root rot, a newly described disease of this foliage plant, 211
Phloem necrosis: of American and slippery elm in New York, disease is in epiphytotic proportions in the Finger Lakes region, 159; of American elm in Pennsylvania, 525
Phoma destructiva: 869
--- exigua var. exigua: 238
--- medicaginis: 286

Phomopsis sp.: 103; 995
--- sclerotioides: 53
Phragmidium mucronatum: 1069
Phyllactinia corylea: 313; 852
Phyllosticta sp.: 11
Phymatotrichum omnivorum: 328
Physalospora obtusa: 1102
Physiological diseases: stem-cavity of McIntosh apples, incidence appears highest in early havested fruit, 543; late-spring leaf scorch of maple and beech trees, 526
Physiological vigor: as a basis for selection for crown resistance to Dutch elm disease, 725
Phytophthora spp.: 927
--- ? arecae: 678
--- cactorum: 108
--- cinnamomi: 290
--- citrophthora: 927
--- megasperma: 472; 830
--- --- var. sojae: 536
--- nicotianae var. nicotianae: 678
--- --- var. parasitica: 678; 689
--- palmivora: 71
--- parasitica: 927; 971
--- --- var. nicotianae: 593
--- phaseoli: 65; 237
Pigment glands: in cotton, as a diagnostic test of the pathogenic properties of new isolates of Xanthomonas malvacearum, 765
Pink root: of onion, caused by Pyrenochaeta terrestris, occurs naturally in Oregon soil, 180
Pinus albicaulis (whitebark pine): Bifusella saccata reported for first time in California on Pinus albicaulis, 102
--- ayacahuite (Mexican white pine): first report of Coleosporium crowellii, a microcyclid rust parasite on Mexican white pine in southern and northern Oaxaca, 474
--- cembroides (piñon pine): new host of Coleosporium crowellii in Mexico, 474
--- clausa (sand pine): no resistance to lance nematode found in two races of sand pine, 691
--- contorta (lodgepole pine): Arceuthobium americanum, parasitism of mistletoe to this pine is closely related to the altitude in Colorado, 223
--- cooperi: rust infection of limbs appears to be Peridermium filamentosum, in Mexico, 896
--- culminicola: new host of Coleosporium crowellii in Mexico, 474
--- divaricata (jack pine): infected with Scleroderris canker, 245
--- durangensis: affected with pine rust, in Mexico, 896
--- echinata (shortleaf pine): air pollution caused some damage to trees growing 1000 yards from source, 3
--- edulis (piñon pine): new host of Coleosporium crowellii in southern Utah, 474
--- elliottii (slash pine): pine seedlings artificially inoculated showed an apparent anatomical mechanism for gall-rust resistance, 531; an improved method for large-scale inoculations with Cronartium fusiforme, 931
--- flexilis (limber pine): Arceuthobium cyanocarpum, its distribution and limitations in the Colorado Front Range, 223
--- jeffreyi (Jeffrey pine): control of Sirococcus tip blight with chlorothalonil, Difolatan and zinc + maneb fungicides, 241
--- michoacana: limb rust, in northern Mexico, 896
--- monophylla: new host of Coleosporium crowellii in southern California, 474
--- nigra austriaca (Austrian pine) first record of Scleroderris lagerbergii on this host in North America, 245
--- ponderosa (ponderosa pine): Arceuthobium vaginatum, a host-parasite combination in Colorado where distinct altitudinal zones of parasitism exist, 223; a strong relation exists between bark beetles and root rots in Idaho, 498; pine limb rust fungi in Mexico, 896
--- resinosa (red pine): host of Scleroderris lagerbergii in New Brunswick, Canada, 245; a new host for brown spot needle fungus, Scirrhia acicola in Wisconsin, 712
--- strobus (eastern white pine): acute symptom expression from exposure to oxides of nitrogen and sulfur dioxide, 3; Scleroderris lagerbergii infection, 245
--- sylvestris Scotch pine): brown spot needle blight on Christmas trees in Wisconsin, Bordeaux mixture gave good control, 99; Scleroderris canker infection in New Brunswick, Canada, 245; incidence of Lophodermium needle cast in North Carolina State Forest Nursery and control with maneb, 457; severely galled Christmas tree infected with Endocronartium harknessii found near Bloomsburg, Pennsylvania, 1058
--- taeda (loblolly pine): chronic damage caused by nearby air pollutants in Virginia, 3; stump inoculation with Peniophora gigantea against Fomes annosus, 66; four fungicides have little if any systemic activity against the fusiform rust fungus, 628; inoculation method that produces uniform infections of Cronartium fusiforme on pine and oak hosts in testing program for fusiform rust resistance, 931; further tests in Virginia with chain saw-applied

Peniophora gigantea in loblolly pine stump inoculation for control of Fomes annosus, 958
--- virginiana (Virginia pine): chlorotic mottle caused by oxides of nitrogen and sulfur dioxide, 3
Piricularia sp. (Pyricularia sp.): 157
--- grisea: 210
--- oryzae: 960
--- setariae: 740
Plant age: of sweetpotatoes in relation to symptom expression of virus infection, 501
Plant Introduction lines: three lines of Pisum sativum show resistance to pea fizzletop disease, 131
Planting media: uptake of thiabendazole in elm seedlings was best in sand or perlite and greatly reduced with peat, 855
Plasmodiophora brassicae: 375
Plasmopara halstedii: 428
Plenodomus destruens: 219
Plum: brown rot, caused by Monilinia fructicola, controlled effectively with piperazine derivative fungicide, 456
Podosphaera leucotricha: 451; 770
Poinsettia: poinsettia wilt in glasshouse-grown plants caused by lime in potting mix, 415; new disease, bacterial stem rot, 480; effects of Botrytis and ozone on bracts and flowers of poinsettia, 814
Poison oak (Rhus toxicodendron): found to be host of pecan pathogen, Cristulariella pyramidalis, 176
Poland: 462
Polyporus hispidus: 300
--- lucidus: 108
--- schweinitzii: 498
--- spraguei: 406
--- sulphureus: 406
Populus spp.: additional hybrid poplar hosts of Fusarium canker in Ohio, 523
Poria weirii: 498
Postharvest decay: of apple, 'Jonathan' apples in cold storage found infected with a dark rot caused by Hormiscium sp., 275
---: of blueberry, Alternaria tenuissima was primary pathogen of highbush blueberry fruit in North Carolina, 118
---: of citrus, by Alternaria citri, during extended storage, 909
---: of lime, in Indian local markets, caused by Aspergillus niger, 64
---: of mango, heat treatments and hot fungicides caused fruit scalding, 751
---: strawberry, preharvest fungicide treatments for control of pre- and postharvest fruit decay of strawberry, 296
---: sweetpotato, surface rot of sweetpotato, caused by Fusarium spp., 783
Potato: occurrence of Ulocladium consortiale as an associate fungus of Alternaria solani in potato tuber blight, 308; three inoculation methods used to screen potatoes for resistance to Rhizoctonia solani evaluated, 348; method for detecting contamination by Erwinia carotovora in tubers, 552
Powdery mildew: tillage and pruning of cultivated plants is related to their susceptibility to infection by powdery mildews, 852
---: of apple: controlled in New York with CELA W 524, 451; eradication of apple powdery mildew from infected buds with pruning agents, 770
---: of cantaloup, caused by Sphaerotheca fuliginea, 935
---: of grasses: benomyl provided longterm control of powdery mildew on orchardgrass, 75; field resistance to Erysiphe graminis in a collection of Poa pratensis (Kentucky bluegrass) mostly from Alaska, 618
---: of cucumber, controlled effectively with triarimol, 206
---: of kalanchoe, controlled with folpet and benomyl, 405
---: of oak, four species occur on 28 species of oak in the San Francisco Bay area in California, 313
---: of squash: controlled with triarimol, 45; of squash and cucumber, controlled effectively with a new systemic fungicide, PP 675, 878
---: of strawberry: control with three systemic fungicides, 235; benomyl in foliar applications controlled infection in Louisiana, 368
---: of wheat, controlled with two fungicides, Morestan or BAY 79770, 358
Pratylenchus coffeae: 626
--- penetrans: 141; 577; 914
Precipitation: relationship between maple canker incidence and precipitation, 942
Proboscidea jussieui: natural occurrence of bean yellow mosaic virus in Proboscidea, 548
Prunus (see also Common names): apple chlorotic leaf spot virus, possibly responsible for the bark split disease of prune trees, 293; peach and almond seedlings translocated oxycarboxin more readily when they were pretreated with two growth retardants, 590; root chip inoculation, an indexing method for Prunus stem pitting, 785
Pseudomonas syringae: 325; 490
--- stizolobii: 446
Pseudopeziza medicaginis: 286
Pseudotsuga menziesii (Douglas-fir): widespread incidence of Rhizina root rot

found to be associated with Douglas-fir seedlings planted on burned-sites in Washington, 407; significant associations between bark beetles and root rots of Douglas fir found in Idaho study, 498; a canker disease caused by Dermea pseudotsugae found on young Douglas-firs in California, 843
Pterocarpus angolensis (mukwa): mukwa blight in Central Africa is causing a dying out of this valuable tree species, 820
Puccinia arachidis: 290; 382; 873
--- chrysanthemi: 939
--- chondrillina: 858
--- coronata var. coronata: 614
--- graminis f. sp. avenae: 608
--- --- f. sp. tritici: 410; 554; 1038
--- pelargonii-zonalis: 759
--- recondita f. sp. tritici: 77; 410; 534; 827
--- stenotaphri: 630
--- striiformis: 1071
--- zoysiae: 227
Puerto Rico: 648
Pyrenochaeta terrestris: 180
Pyricularia sp. (Piricularia sp.): 157
--- grisea: 210
--- oryzae: 960
--- setariae: 740
Pythium spp.: 328
--- aphanidermatum: 507; 840; 1043
--- debaryanum: 973
--- mamillatum: 973
--- myriotylum: 600
--- splendens: 211; 973; 1074
--- ultimum: 973

Quercus (see Oak)

Races: new race of anthracnose on kenaf is apparently present in Zambia, 1049
---: race 1 of Fusarial wilt of banana, caused by Fusarium oxysporum f. sp. cubense responsible for disease attack on 'Dwarf Cavendish' in the Canary Islands, 1000
---: race T of Helminthosporium maydis: varying levels of resistance to race T in corn hybrids were attributed to nuclear genes interacting with cytoplasmic factor(s), 112; production and relative host-specificity of a toxin from race T, 247; live plant detection of airborne spores, 659; in Iowa in 1971, 660; races O and T of H. maydis produce toxins, that are nonspecific and host-specific, respectively, 805; race O of H. maydis on a corn hybrid in normal cytoplasm has greater ability to be the dominant species over race T on the same hybrid, 889
---: new race of Phytophthora megasperma var. sojae is pathogenic to soybean, 536
---: race 1 of Phytophthora parasitica var. nicotianae found in east Tennessee, 593
---: Puccinia coronata var. coronata, crown rust, Goar and Ky 31 races were found on tall fescuegrass in Alabama, 614
---: of Puccinia graminis f. sp. avenae in the USA during 1971, 608
---: Puccinia graminis f. sp. tritici: wheat stem rust in the United States in 1971, the three most predominant races were 15B-2, 11-32-113, and 151, 554; wheat stem rust races in the Yaqui Valley of Mexico in 1972, 1038
---: Uromyces phaseoli, new race of bean rust isolated from pole beans in Florida, 759
Radiation: on lemons to prolong storage life caused fruit injury, 582
Radioautography: of almond, chestnut and peach seedlings, revealed enhanced penetration and translocation of oxycarboxin following foliar treatments with SADH and CCC growth retardants, 590
Radopholus similis: 85
Ramularia sp.: 328
Rape (Brassica campestris) (see Crucifers)
Raspberry: red, fumigation of soil decreased root rot of raspberry but residual effects were reduced after 1-4 years, 467; calico, a transmissible disorder of red raspberries in Washington, 779; 65 commercial red raspberry fields in Washington were indexed for virus infection, 23% showed infection but tomato ringspot virus was the only virus causing visible damage in the field, 1007
Rate: of pathogenic and saprophytic development of Fomes annosus in roots of dominant and suppressed eastern redcedar, 987
Red ring disease: of coconut palm, caused by nematode Rhadinaphelenchus cocophilus, in Grenada, West Indies, 339
Relative humidity: importance in the potential for spread and increase of southern corn leaf blight, 104
Resistance: to zoysia rust, zoysia 'Emerald' and 'Beltsville 101' had best ratings for rust resistance of 14 selections tested, 227; of lima bean lines and selections to Rhizoctonia solani, 268; anthracnose resistance in F_2 populations of 10 tomato introductions shows no clear segregation ratios of resistant to susceptible, 321; to Ditylenchus dipsaci in onion species and hybrids and backcross progeny, 329; of potato varieties to Rhizoctonia solani,

348; to peanut rust in peanut varieties and accessions, 382; of Xanthomonas dieffenbachiae isolates to streptomycin, 394; sunflower inbred line HA 61 showed high resistance to downy mildew in field evaluation, in North Dakota, 428; anatomical resistance to gall-rust infection in slash pine, 531; monogenic resistance and polygenic tolerance approaches for host:pathogen interaction are described and discussed in relation to controlling Fusarium wilt of tomato with resistant varieties, 695; of lettuce lines to Sclerotinia sclerotiorum, 875; many spring barley cultivars showed immunity to Helminthosporium stripe in five inoculation tests with eight isolates of H. gramineum, 891; of Citrus x Poncirus hybrids to feeder root infection by Phytophthora spp., 927
Rhadinaphelenchus cocophilus: 339
Rhizina undulata: 407
Rhizoctonia sp.: 990
--- solani: 268; 328; 348; 400; 511; 840; 1054
Rhizopus nodosus: 990
Rhopalosiphum maidis: 885
Rhubarb (Rheum rhaponticum): first record of rhubarb as a host for Meloidogyne hapla, 417
Rhynchosporium oryzae: 106
Rhyncophorus palmarum: 339
Rhytisma acerinum: 108
Rice: Helminthosporium rostratum found on rice disease specimens from Myrtle Beach, South Carolina, a first report for the United States, 26; leaf scald, caused by Rhynchosporium oryzae, in West Africa, 106; insecticide PP-211 when added to fungicide-treated rice seed improved control of seedling rot and diseases, 154; chemical applications of nematicides in rice fields found to stimulate competitive grassy weeds, 420; blast, virulent and less virulent races of the blast pathogen have the same ability to germinate and form appressoria on rice leaves, 960
Root and butt decay: of oak trees in ferneries caused by cultivation practices, 811
Root chip inoculation: indexing method for Prunus stem pitting, 785
Root crops: and associated plant-parasitic nematodes in the West Indies, 1077
Root diseases: of apple trees, in New York, preplant treatments with nematicides increased growth of apple trees and initially reduced populations of Pratylenchus penetrans, 577
Rose: rose rust, caused by Phragmidium mucronatum, not controlled with folpet, 1069
Rot: black rot, of sweetpotato, thiabendazole prevented development on inoculated sweetpotatoes, 219
---: boll rot: of cotton: caused by Aspergillus flavus, its geographical distribution in experimental plots as revealed by fiber fluorescence, 215; estimated reduction in cotton yield from boll rots in 1971, 328
---: brown rot of plum, controlled with piperazine derivative fungicide, in New York, 456
---: charcoal rot of cucurbits, isolated from five cucurbit hosts, 115
---: collar, of koa (Acacia koa) in Hawaii, 73
---: crown and root rot of forage crops caused by Sclerotinia trifoliorum as affected by north- and south-facing slopes, 371
---: crown of wheat, seedling blight, root and crown rot disease complex, 788
---: crown and leaf and scape spot of statice, in Florida, 894
---: flower, of camellia, in Alabama, 919
---: foot rot of sweetpotato, thiabendazole greatly reduced incidence, 219
---: ---: of wheat: caused by Cercosporella herpotrichoides, controlled with benomyl, 20; Cercosporella foot rot of winter wheat, controlled with benomyl, 178; control with benomyl dependent on the rate and time of application, 342; water relations of C. herpotrichoides studied in vitro indicate the pathogen is active within the wheat stem at temperature and moisture levels previously considered limiting to it, 594
---: fruit: of lime, caused by Aspergillus niger in local Indian markets, 64; of highbush blueberry in North Carolina, 118; of apple, by Hormiscium sp., 275; of strawberry, preharvest fungicide treatments to control fruit decay, 296; dry rot of guava fruits in the orchard in India, caused by Diplodia natalensis, 323; of mango, stem-end rot was not controlled with fungicides or hot water treatments, 751
---: pod rot: of peanut, soil management practices in Israel in relation to Pythium infection, 600; pod and collar rot of soybean, caused by Fusarium semitectum, 693
---: Rhizoctonia hypocotyl rot of lima bean, 'Jackson Wonder' and S-400 showed more tolerance to R. solani than other varieties in greenhouse and field tests, 268
---: root: black root rot of bean and tobacco controlled with several fungicides, 15; controlled in tobacco seedbeds with fungicides in New Zealand, 708

---: ---: of centipedegrass, caused by Fusarium roseum, 42
---: ---: of conifers in Idaho, associated with bark beetle damage, 498
---: ---: of cucumber, caused by Phomopsis sclerotioides, 53
---: ---: Fomes annosus: stump treatment for biological control, 66; 958
---: ---: Phymatotrichum omnivorum, loss estimate of cotton crop for 1971, 328
---: ---: Phytophthora: of raspberry, soil fumigation improved yield and growth performance over check, 467; of alfalfa, caused serious losses in Arizona, 472; of alfalfa, caused by P. megasperma, in Washington, 830
---: ---: Pythium: previously undescribed disease of Philodendron panduraeforme and Scindapsus aureus, caused by P. splendens, found in Florida, 211; P. aphanidermatum found to possess the full complement of survival mechanisms to persist under adverse climatic conditions and to be widely distributed in Arizona, 507; P. splendens caused severe stunting of cotton plants in greenhouse, but severe root rots were not observed under field conditions in California, 973
---: stalk, of corn, fungi associated with corn stalk rot in Pennsylvania in 1970 and 1971, 836
---: stem, brown stem rot of soybean, a defoliating isolate of Cephalosporium gregatum significantly reduced yields of inoculated soybeans, 580
---: storage, surface rot of sweetpotato, caused by Fusarium spp. after storage, cultivars show wide and consistent differences in incidence, 783
---: tuber rot of Dioscorea floribunda (a steroid sapogenin-containing species) controlled with benomyl, 990
Rotylenchulus reniformis: 551; 1040; 1077
Rotylenchus sp.: 1059
Rubber plant (Ficus elastica): relationship of disease to methods of inoculation of F. elastica with Aphelenchoides besseyi, 471; artificial and natural infection with Cercospora disease, 1091
Rust: four cultures of rust fungi can be applied simultaneously to single wheat seedling with multiculture inoculation system, 847
---: of apple, controlled in New York with piperazine derivative fungicide, CELA W 524, 451
---: of pole bean, a new race isolated in Florida is designated race 35, 759
---: crown rust of tall fescue, two races of the pathogen found in Alabama, 614
---: flax, systemic protectant and eradicant chemical control with oxycarboxin or U8342, 737
---: gall: anatomical resistance to infection in slash pine, 531; fusiform, of loblolly pine, selected systemic fungicides provided little control of fusiform rust of loblolly pine in forest tree nurseries, 628; western gall rust of hard pines found in Pennsylvania, 1058
---: geranium, found on geranium cuttings in Pennsylvania, 759
---: of kenaf, varietal resistance to Aecidium garkeanum, in Zambia, 1047
---: leaf, of chrysanthemum, 52 cultivars tested for susceptibility in 1969, and 15 in 1972, 939
---: ---: of wheat: distribution of virulence of Puccinia recondita f. sp. tritici in North Dakota in 1971, 77; in Georgia, the 1971-1972 wheat crop threatened by severe leaf rust infection, 534; a culture of Puccinia recondita virulent to 'Transfer' wheat was isolated in Indiana, 827
---: limb rust (systemic crown kill) of pine, in Mexico, 896
---: of peanut: rust was present in 24 counties in Georgia in late summer, 1971, the most widespread distribution on record, 290; five PI introductions and one breeding line, NC 13, showed marked resistance to peanut rust in tests in Jamaica, 382; caused economic losses in early crop of Spanish peanuts in south Texas for the first time in 1971, 873
---: St. Augustinegrass in the Gulf Coast area of Texas infected with Puccinia stenotaphri, 630
---: stem: of oat, prevalence of races in the U.S. in 1971, 608
---: ---: of wheat: races of Puccinia graminis f. sp. tritici in the USA during 1971, 554; in the winter nursery in the Yaqui Valley, Mexico, 1038
---: stripe, of wheat, the major factors affecting the amount of yield reduction from stripe rust in wheat in northeast Oregon in 1971, 1071
---: zoysia, conditions affecting the development and perpetuation of the rust, 'Emerald' and 'Beltsville 101' had best ratings for rust resistance, 227

Safflower (Carthamus tinctorius): Frio was most tolerant cultivar of 12 tested for sensitivity to ozone, 195
Salt injury: effect of salt injury on shoot growth of sugar maple in New Hampshire, 1004
Sanitation: for control of anthracnose of strawberry by plant selection, 515
Sassafras albidum: host of zonate leafspot of

pecan, 176
Scab: of apple: infections on twigs of more species and varieties of flowering crab apple, 252; prevention of primary scab infection in an isolated apple orchard 1970-71, 736; late season application of benomyl + surfactant F to apple trees in Chile gave almost complete inhibition of the perfect stage of Venturia inaequalis, 1079
---: of pecan, controlled in Oklahoma with benomyl sprays, 980
Schizaphis graminum: 885
Scindapsus aureus: discription and control of Pythium root rot, a newly described disease of this plant, 211
Scirrhia acicola: 99; 108; 712
Scleroderris lagerbergii: 245
Sclerophthora macrospora: 773
Sclerospora sorghi: 740
Sclerotinia sclerotiorum: 798; 875
--- trifoliorum: 371
Sclerotium rolfsii: 362; 656
Scotland: 803
Scurf: of sweetpotato: heated dips of benomyl and thiabendazole were highly effective for control of scurf, 122; thiabendazole eliminated scurf in greenhouse pot tests, 219
Seed: interaction among microorganisms occurring naturally and applied to pericarps of corn kernels, 861
---: storage: of sunflower seeds, 6.5% moisture content, or lower, will prevent invasion by storage fungi, 173; of popcorn, a combination of a moisture content below 14.0% and a temperature of 10-20°C should be suitable for long-time storage without loss in quality from storage fungi, 1097
---: transmission, seed-transmitted bacterial diseases in Michigan navy (pea) beans, 378
---: treatment: of bean with benomyl significantly reduced root rot caused by Thielaviopsis basicola, 15; fungicides + insecticide PP-211 on rice seed improved control of disease over fungicides alone, 154; several fungicides used as seed treatments reduced soilborne Helminthosporium stripe of barley, 183; for control of Pythium blight of ryegrass, 1043
Seedling: growth stunting of eastern white pine caused by air pollution, 3; peach, longest survival of root-knot nematodes on 'Nemaguard' seedlings was at constant temperature of 30°C, 305; 27 spring barley cultivars showed immunity and 13 showed resistance to barley stripe in inoculation tests, 891; resistance to feeder root infection in Citrus x Poncirus hybrids, 927; wilt symptoms were restricted in tomato seedlings grown in soil treated with thiabendazole following stem inoculation with Fusarium oxysporum f. sp. lycopersici and Verticillium albo-atrum, 1054
Seedling diseases: seedling tip blight of Arizona Cypress and eastern red cedar, caused by Curvularia intermedius, 192; Sirococcus tip blight of Jeffrey pine seedlings controlled with fungicides, 241; citrus, grafted and ungrafted 1- and 2-year-old nursery seedlings were affected by disease complex caused by Fusarium sp. and a nematode, in Tanzania, 318; loss of cotton in U.S. by State from seedling diseases, 328; Douglas fir, seedling mortality associated with Rhizina root rot in burned and planted clear-cuts in western Washington, 407; needle cast of Scotch pine controlled with maneb, 457; male cytoplasm as well as female cytoplasm affects the reaction of corn seedlings to seedling disease (Fusarium sp.), 575; resistance of hard red spring wheat and durum wheat to seedling blight and crown rot, 788
Septoria nodorum: 188; 761
--- tritici: 983
Serology: of lilac mottle isolated from Maryland lilac bush, 923
Shade: shade and wet soil were found to increase southern corn leaf blight in a field, 203
Shoot-blight disease: of western hemlock in Alaska and British Columbia, 645
Shuck disease: of pecan, one form, stem-end blight, reduced with prepollination sprays of benomyl and TPTH, 276
Sirococcus strobilinus: 241; 645
Skeleton weed (Chondrilla juncea): Puccinia chondrillina as a biological control agent will infect and sporulate over a wide temperature range, 858
Slope direction: amount of disease incidence in five forage species was influenced by the direction of the slope, that is, facing north or south, 371
Slope preference: and infection severity of dwarf mistletoe species in Boulder County, Colorado, distinct altitudinal zones of parasitism specific for each host-parasite combination exist, 223
Smut: flag: of Poa pratensis 'Merion', controlled with EL-279, 55; controlled with BAS 3191 F, 252; controlled on wheat, 636
---: stripe: of Poa pratensis 'Merion', controlled with EL-279, 55; eradicated with BAS 3191 F, 252; controlled with three systemic fungicides, 334

Snow mold: resistance of 56 bluegrass cultivars to Typhula itoana and Fusarium nivale evaluated in Michigan, 32; Fusarium nivale is cause of a leaf blight on wheat in Scotland and on triticale in Mexico, 803
Soil: pH, incidence of Verticillium wilt increased and Fusarium wilt decreased on tomato when soil pH was raised from 6.0 to 7.0 or 7.5, 151; Pyrenochaeta terrestris, the onion pink root fungus, is naturally occurring component of soil in Oregon, 180; influence of wet soil and shade on southern corn leaf blight, 203; mean incidence of Fusarium canker of yellow poplar in South Carolina was positively correlated with increasing levels of potassium in the soil, 310; survival of Pythium aphanidermatum under arid soil conditions in Arizona, 507; sodium hexametaphosphate to disperse clay particles was most effective method in extracting nematodes from clay soils, 808
---: amendments: Alternaria leaf spot more prevalent on benomyl-treated carnation plants than on untreated controls, 9; alfalfa hay and castor pomace partially effective against black root rot of bean and tobacco, 15; effect of micronutrients on Verticillium and Fusarium wilt of tomato, 151; fungicidal soil drenches + mulch reduced stem blight of new plants of Vinca minor, 238; reduction of ozone injury of greenhouse-grown bean seedlings with triarimol and EL-279 as soil amendments, 386
---: treatments: EL-279 effective against stripe smut and flag smut of bluegrass, 55; complete fertilizer, preplant nematicide treatments, brush burning, and fertilizer + nematicide treatments increased yields and growth of sour cherry trees, 141
---: ---: for fungi: soil drench application of benomyl gave control of powdery mildew on orchardgrass, 75; peaches in fumigated soil had less stem pitting than peaches in nonfumigated plantings, 162; benomyl and thiabendazole as soil drenches controlled stem blight of Vinca minor, 238; BAS 3191F in single application eliminated flag and stripe smuts from 'Merion' bluegrass, 252; potassium azide reduced symptoms of Rhizoctonia solani and Verticillium wilt in potatoes, 400; soil fumigation and its effect on raspberry root rot, in Washington, 467; use of steam-sterilized soil has almost completely eliminated cyclamen wilt from California nurseries that used to suffer total loss of their crops, 493; fungicides to control Dutch elm disease, 604; soil drench of oxycarboxin or U8342 effective against flax rust, 737; several fungicides tested under grower conditions for control of Pythium splendens on Scindapsus aureus, in Florida, 1074; drench treatments with six fungicides for control of stripe smut of Kentucky bluegrass, 1083
---: ---: for nematodes: preplant soil fumigation with ethylene dibromide or dibromochloropropane decreased root-knot index, and peach trees showed increased fruit yield, 169; D-1410 mixed with soil, controlled tobacco cyst nematode, 255; benomyl drench suppressed Heterodera avenae on barley, 261; potassium azide ineffective against nematodes when applied after emergence of peanuts, 362; with low vapor pressure nematicides to control sugarbeet nematodes on Brussels sprouts, 375; apple trees showed increased growth after preplant treatment with nematicides, 577; two cycles of flooding followed by drying for 2 weeks reduced nematodes in celery seedbeds on muck soil in Florida, 661; soil fumigation with 1,3-D and methyl bromide controlled Xiphinema index and fanleaf virus, 1031
Sorghum: cytoplasmic vulnerability to Helminthosporium maydis is not apparent in sorghum cultivars, 682; disease survey of sorghum and millets in India, 740; downy mildew was first observed in Arizona in 1971 on wheat, barley, triticale and sorghum, 773
South Carolina: 26; 169; 310; 1087
South Dakota: 572; 898
Southern corn leaf blight: 104; 112; 203; 485; 659; 660; 682; 767; 866; 889; 969; 1087
Soybean: Endogone sp. found in soybeans and sweetpotatoes in Louisiana for the first time, 58; peanut mottle virus found in commercial soybeans in Georgia, 146; root rot, a new race of Phytophthora megasperma var. sojae found that is pathogenic to soybean cultivars, 536; brown stem rot, soybean plants inoculated with Cephalosporium gregatum defoliating isolates showed reduced yields, 580; pod rot and collar rot, caused by Fusarium semitectum in India, 693; five soybean varieties tested for lateral root production and to galling after infection with Meloidogyne javanica and M. incognita, 882
Sphaeropsis ulmicola: 108
Sphaerotheca fuliginea: 206; 852; 935
--- humuli: 368; 405
--- lanestris: 313

--- macularis: 235
Spinach: naturally infected by lettuce mosaic virus in New York, no tolerance or resistance found in 24 spinach cultivars, 281; cultivars Bounty and Winter Bloomsdale were least sensitive of six cultivars tested for ozone sensitivity in Massachusetts, 832
Spinacia oleracea (see Spinach)
Spores: ascospores as the primary infective propagule of Sclerotinia sclerotiorum on lettuce in Wisconsin, 798; a method is given for calculating the sampling efficiency of rotorod samplers, 704; live plant detection of airborne spores of Helminthosporium maydis race T in the field, 659
Spore germination: a method for collecting ascospores of Venturia inaequalis for quantitative studies on spore germination, 39; no significant differences in spore germination of Fusarium solani were observed in extracts from cankered and uncankered sugar maple trees, 944
Spore traps: cylindrical rods wrapped with a polyethylene square coated with Dow-Corning 200 fluid have higher trapping efficiency than flat slides coated with adhesive, 719
Sporulation: of Scirrhia acicola, that causes brown spot needle blight on Scotch pine Christmas trees in Wisconsin, 99
Spring black stem: of alfalfa, controlled with fungicides, 286
Squash (see Cucurbits)
Statice (Limonium spp.) Sclerotium rolfsii is pathogen of statice in Florida, possible first report, 656; L. perezii, four aphid species transmitted turnip mosaic virus from statice to statice in California fields, 722; L. sinuatum, a new crown rot, leaf and scape spot is incited by Colletotrichum sp. in Florida, 894
Stem-cavity browning: of McIntosh apples, 543
Stem-end rot: of mango, fungicidal or hot water treatment did not control decay, 751
Stem tissue extracts: of sugar maple, and their influence on spore germination of Fusarium solani, 944
Stemphylium botryosum: 286
--- loti: 371
Stem pitting: of peach, in fumigated and nonfumigated field plantings in New Jersey, 162; of sour cherry in East Europe, 462; of peach, in New York, 616; indexing by root chip inoculation was reliable and feasible for selecting peach propagating material free from stem pitting, 785
Stereum frustulatum: 406
Storage: moisture content of sunflower seeds in relation to invasion by storage fungi, 173; sound clean popcorn seed, even of fairly high moisture content, can probably be kept for some time without damage from storage fungi if held at a constantly low temperature, 1097
Storage life: of lemon, irradiation controlled Geotrichum candidum decay, but injured fruit of lemon, 582
Strains: of tobacco etch virus and potato virus Y that cause severe losses to pepper in Florida, 586; strain E of sugarcane mosaic virus occurred in St. Augustinegrass in south Florida, 822; strain E of sugarcane mosaic virus from St. Augustinegrass, its characterization, 885
Straw breaker foot rot: of winter wheat, 594
Strawberry: leaf blight, caused by Dendrophoma obscurans, controlled with benomyl or thiabendazole in Florida summer nursery, 23; preharvest fungicide treatments increased yields over untreated controls, 296; temperatures between 65 and 75°F appear to be optimum for fungus growth of leaf spot organism, Mycosphaerella fragariae, 345; powdery mildew of strawberry controlled with benomyl in Louisiana, 368; anthracnose, control of crown rot phase in Louisiana by sanitation, plant breeding and benomyl spray, 515; cultivars show differences in tolerance to virus diseases in the Pacific Northwest, 792; occurrence of strawberry leaf blotch in Florida, 998
Stump treatment: of loblolly pine, Peniophora gigantea applied by chain saw against Fomes annosus, 66; 958
"Sudden death": of citrus trees on Poncirus trifoliata rootstock, New South Wales, Australia, 80
Sugarbeet: influence of inoculation date on expression of beet western yellows (BWYV) tolerance in 12 varieties, 93; 26 isolates of BWYV tested to determine effects of virus on growth of sugarbeet varieties, 198; triphenyltin chloride showed systemic activity in sugarbeet seedlings and inhibited growth of Alternaria tenuis in laboratory tests, 776
Sunflower (Helianthus annuus): moisture content of sunflower seeds in relation to invasion by storage fungi, 173; field evaluation of sunflower for downy mildew resistance, 428
Survey: of viruses infecting tomato in Louisiana, 272; of air pollution damage to vegetation in Pennsylvania in 1969, 701

Survival: of Meloidogyne hapla on rhubarb roots served as infection for tobacco plants inside greenhouse in Ontario, 417; occurrence and survival of Pythium aphanidermatum under arid soil conditions in Arizona, 507; relative survival of populations of race O and T of Helminthosporium maydis on a corn hybrid in normal cytoplasm, 889; of H. maydis under field conditions in South Carolina, 1087
Susceptibility: of some tomato varieties and inbred lines to bacterial leaf spot, 504; of chrysanthemums to leaf rust in Florida, 939; various cultivars of lettuce appear to differ in susceptibility to injury from increased concentrations of CO_2, a normal respiratory product of lettuce, 1051
Sweetpotato: first report of an Endogone sp. from roots of sweetpotatoes and soybeans in Louisiana, 58; an internal disorder of unknown cause yields hard inedible tissues when roots are cooked, 88; control of scurf with fungicide dips of benomyl or thiabendazole, 122; fungicidal treatment of sweetpotato with thiabendazole prevented black rot, eliminated scurf, and reduced foot rot in greenhouse pot tests, 219; effects of plant age, light intensity and leaf pigments on symptomatology of virus-infected sweetpotatoes, 501; clonal differences in the incidence of surface rot (Fusarium spp.) suggest that a genetic basis for resistance exists, 783
Sycamore (Platanus occidentalis): fungicidal control of the twig blight stage of sycamore anthracnose, 354
Symptom expression: of eastern white pine to air pollutants, 3

Tanzania: 318; 678
Target spot: of tomato, disease of tomato new to the United States, 243
Taro (see Colocasia esculenta)
Tecomaria capensis: a previously unreported host of alfalfa mosaic virus, in Israel, 1035
Temperature: heated dips (49-55°C) of benomyl or thiabendazole controlled sweetpotato scurf, 122; hot water treatments, dasheen mosaic virus not inactivated in Xanthosoma corms below 65°C, and above that temp the corms did not germinate, 320; mango fruits held at 55°F for 1 week after hot water or heated fungicide treatment developed scald injury, but not at 70°, 751; moisture and temperature requirements for storage of popcorn, 1097
---: air: Madrone tree canker infection found to be most rapid at 25°C, 50; temperature, dew period, relative humidity and air flow requirements for potential spread and increase of southern corn leaf blight, 104; effect of temperature and light on colonization and sporulation of Botrytis cinerea on geranium, 126; influence of temperature on infection of Meloidogyne incognita acrita on 'Nemaguard' peach seedlings, 305; effect of different temperatures on growth of Mycosphaerella fragariae on strawberry leaves, 345; at 20°C 'Monroe' bean inoculated with bean common mosaic virus (BCMV) consistently produced local lesions, at 16° and 28° lesion number and quality were poor, 714; effect of dew temperature, post-dew temperature and dew period on infection of a male-sterile corn hybrid by race T of Helminthosporium maydis, 767; effects of temperature and duration of leaf wetness on infectivity of Puccinia chonrillina to skeleton weed, 858; abnormally low temperatures and drought conditions in Iowa resulted in light to moderate development of southern corn leaf blight in Iowa in 1971, 866; several viruses of citrus were inactivated with alternating temperatures day-night at 40/30°C, 976; temperature and rainfall influence incidence of Curvularia corm disease in Illinois, 995
Tennessee: 593
Texas: 45; 300; 630; 873; 1065
Theobroma cacao (cocoa): cocoa swollen shoot virus isolate from Nigeria appears to be unique, 755; cocoa mealypod disease is more common in Ghana than previously indicated, 1011
Thermotherapy: of citrus for inactivation of certain viruses, 976
Thielaviopsis basicola: 15; 708
Tikka leaf spot (Cercospora): of peanut, controlled in India with benomyl, 1099
Tobacco: leaf injury induced by coincident applications of a sucker control agent and a fungicide, 338; infection of greenhouse-grown tobacco by Meloidogyne hapla found to be caused by rhubarb roots growing under greenhouse footing from outside, 417; black shank, race 1 of the black shank pathogen found in Tennessee, 593; control of black root rot of tobacco with benomyl and thiophanate-methyl, 708
Tomato: the economic impact of tomato diseases in Montserrat, West Indies, 817; translocation, disease remission and persistence of thiabendazole and ETMT in tomato seedlings, 1054
---: anthracnose, the resistance in F_2 populations derived from 10 tomato

introductions, 321
---: bacterial canker, control in hydroponically grown plants, with formalin, 139
---: bacterial leaf spot, differences in susceptibility of 14 tomato varieties and lines, 504
---: damping-off, control of damping-off of transplants with Difolatan or zinc + maneb as a dip treatment, 840
---: Fusarium wilt: controlling Fusarium wilt of tomato with resistant varieties, 695; efficiency of DD-MENCS for Fusarium control increased with paper or polyethylene as mulch material, 953
---: leaf blight of tomato, caused by Phoma destructiva in Ascension Island, 869
---: target spot, caused by Corynespora cassiicola, 243
---: Verticillium wilt, incidence of Verticillium wilt increased and Fusarium wilt decreased on tomato by raising the soil pH from 6.0 to 7.0 or 7.5, 151
---: virus: identification of viruses of tomatoes in North Carolina, 92; virus diseases affecting tomato in Louisiana, 272
Toxin: susceptibility of corn lines to Helminthosporium maydis toxin, 29; H. maydis race T toxin can distinguish reliably between corn lines with normal (resistant) cytoplasm and lines with Tms (susceptible) cytoplasm, 247; of H. maydis, and reaction of corn seedlings with different cytoplasms, 432; toxins isolated from culture filtrates of race O and T of H. maydis, 805
Trachysphaera fructigena: 1011
Transmission: raspberry calico, a transmissible disorder, 779; strain E of sugarcane mosaic virus from St. Augustinegrass was transmitted in stylet-borne manner from corn to corn by Schizaphis graminum and Rhopalosiphum maidis, 885; virus transmission from Chenopodium amaranticolor, 1085
Trichoderma sp.: 661
Trichodorus sp.: 661
Trinidad: 551; 618
Trumpet vine (Campsis radicans): host of zonate leafspot of pecan, 176
Tsuga heterophylla (western hemlock): Sirococcus strobilinus found to be cause of shoot blight of western hemlock in British Columbia and Alaska, 645
Tuber blight: of potato, Ulocladium consortiale is apparently a secondary organism living primarily in tuber tissue invaded by Alternaria solani, 308
Turkey: 352
Tylenchorhynchus sp.: 661; 1059; 1077
--- claytoni: 476; 540
--- dubius: 546
--- nudus: 898
Tylenchulus semipenetrans: 186; 747
Tylenchus sp.: 898; 1059
Typhula itoana: 32

Uganda: 901
Ulocladium consortiale: 308
Uredinales: Coleosporium crowellii, a rust parasite found on several pine species in the Western States and Mexico, 474
Urocystis agropyri: 55; 252; 636; 1083
Uromyces phaseoli var. typica: 759
Ustilago striiformis: 55; 252
--- --- var. poae: 334

Vectors associated with: red ring disease of coconut palm, Rhyncophorus palmarum, 339; pearl millet, Cicadulina mbila, 424; turnip mosaic virus, Myzus persicae, 441; Alomae and Bobone diseases of Colocasia esculenta, by Myzus persicae, 597; leafhopper-transmitted disease of corn, Graminella nigrifrons, 652; new virus of pepper in Arizona (AzPV) by Myzus persicae, 731; strain E of sugarcane mosaic virus, Schizaphis graminum and Rhopalosiphum maidis, 885; canker disease of Monterey cypress, Laspeyresia moth may be an effective vector, 992
Vein spot: of pecan, significant correlation found between stem-end blight and vein spot of pecan in Louisiana tests, 276
Venturia inaequalis: 39; 252; 736; 1079
Venus-Fly-Trap (see Dionaea muscipula)
Verticillium sp.: 108; 400
--- albo-atrum: 108; 264; 328; 1054
--- dahliae: 264
--- nigrescens: 264
Vinca minor: control of stem blight with benomyl or thiabendazole as soil drenches, 238
Virginia: 3; 66; 108; 958
Virginia creeper (Parthenocissus cinquefolia): a host of zonate leafspot of pecan, 176
Virus: of tomato, in North Carolina, 92; virus symptoms in sweetpotatoes as affected by plant age, light intensity and leaf pigments, 501; viruslike particles associated with a leafhopper-transmitted disease of corn in Kentucky, 652; Crambe species were susceptible to several viruses but remained symptomless after inoculation, 665; virus tolerance in strawberry cultivars appeared to come from 'Northwest', and clones of Fragaria chiloensis, 792; alternating temperatures (40/30°C) superior to constant temperature for plant survival and possibly for virus inactivation in citrus, 976; viruses in red raspberry in Washington, 1007; addition of K_2HPO_4 + bentonite to Chenopodium amaranticolor inoculum increased infectivity

of inoculum by 7- to 1200-fold, 1085
---: alfalfa mosaic virus: found in Ajuga reptans in New York, a first report, 285; alfalfa mosaic, clover yellow vein, tobacco ringspot, and white clover mosaic viruses were isolated from white clover for the first time in North Carolina, 774; influence of dimethyl sulfoxide on its translocation, 1014; found in unreported host, Tecomaria capensis, 1035
---: Alomae and Bobone, virus diseases of Colocasia esculenta in the British Solomon Islands, described for the first time, 597
---: apple chlorotic leaf spot virus, possible cause of bark split disease of prune, 293
---: bajra (pearl millet) streak, a leafhopper-borne cereal virus in India, 424
---: barley yellow dwarf: in tall fescue, 167; difference in levels of tolerance in oat varieties to two strains of the virus, 230
---: bean common mosaic virus, local-lesion assay of BCMV on 'Monroe' bean, 714
---: bean yellow mosaic virus of Proboscidea jussieui, 548
---: beet western yellows virus (BWYV): BWYV tolerance can be evaluated in sugarbeets although natural spread occurs, 93; variation among Washington isolates, 198
---: of citrus, vein-enation virus on Satsuma mandarins in Turkey, 352
---: cocoa swollen shoot virus (CSSV), one Nigerian isolate differs from other Nigerian isolates studied previously, 755
---: cucumber mosaic virus: caused mosaic disease of Dichondra in California, in 1971, 27; on tomato, 92; response on Chenopodium amaranticolor as affected by the kind of buffer used, 443
---: dasheen mosaic virus, in Xanthosoma spp., 320
---: fanleaf of grape, progress in the control of nematodes by soil fumigation in nematode-fanleaf infected vineyards, 1031
---: lettuce mosaic virus, natural infection of spinach in New York, 281
---: lilac mottle, host range, physical properties, purification and serology, 923
---: maize dwarf mosaic virus: in naturally occurring johnsongrass in Louisiana corn and sugarcane production areas, 622
---: papaya ringspot virus is cause of serious losses of papaya in Hawaii, 331
---: pea fizzletop, three PI introductions appear resistant to virus-induced disease, 131
---: peanut mottle virus, first natural occurrence on soybean in Georgia, 146
---: peanut stunt virus (PSV), first report of PSV in Alabama, 415
---: of pepper: tobacco etch virus and potato virus Y are cause of severe pepper losses in south Florida, 586; a virus of the potato virus Y group is cause of new disease of pepper in Arizona, 731
---: potato virus Y, on tomato, 92
---: St. Augustine Decline virus found for first time in Louisiana, 69
---: sugarcane mosaic virus: strain E was isolated from St. Augustinegrass in south Florida, not recovered from central and northern Florida, 822, particle and inclusion morphology of strain E isolated from St. Augustinegrass in Florida is described for the first time, 885
---: tobacco etch virus, on tomato, 92
---: tobacco mosaic virus, aucuba strain on tomato, 92; in Crambe reaches a high concentration, but plant is symptomless, 665; a fern, Polypodium aureum, is able to support tobacco mosaic virus replication, 1010
---: tobacco ringspot: from naturally infected dogwood, autumn crocus and Forsythia, in Maryland, 336; dimethyl sulfoxide may influence translocation of the virus, 1014
---: tomato: tomato aspermy virus, 92; tomato strain of tobacco mosaic virus, potato virus Y, tobacco etch virus, and cucumber mosaic in Louisiana, 272
---: tomato ringspot virus: in 'Baco Noir' grapevines in New York, 133; the rapid spread in New York, 1062
---: tomato spotted wilt virus, purified from ageratum, 869
---: turnip mosaic virus: isolated from 'Green Ball' cauliflower, 441; turnip mosaic virus-vector relationships in field-grown statice, in California, 722
---: watermelon mosaic: serious disease of summer squash in Georgia, 147; identification of watermelon mosaic virus 1 in El Salvador, 437; watermelon mosaic virus 1, Momordica charantia is most likely source of infection for cucurbits in Florida, 563

Washington: 20; 50; 93; 198; 398; 407; 467; 565; 779; 792; 830; 1007
Washington, D. C.: 725
Watermelon (see Cucurbits)
Wax emulsion sprays: reduced visible ozone injury to Phaseolus vulgaris 'Pinto III', 6
Weather: sudden change from cloudy, wet conditions to bright sunny weather responsible for late-spring leaf scorch of maple and beech trees in Ontario, 526; hurricane Camille damage and its possible role in the persimmon wilt outbreak in Mississippi in 1970, 916
Weeds: nematicide-weed interaction in rice fields where nematicides stimulate grassy weed growth, 420
West Africa: 106
West Indies: 339; 382; 551; 817; 1059; 1077
West Virginia: 371

Wet-rot: of pole beans, a previously undescribed disease caused by Choanephora cucurbitarum, 967
Wheat: hard red spring wheats and durum wheats showed differences in response to seedling blight and crown rot, 788
---: Cercosporella foot rot: controlled in Washington with single benomyl spray, 20; controlled with benomyl at three seeding dates in Oregon, 178; spring application of benomyl + weed control gave better control than in fall, 342; some water relations of C. herpotrichoides, 594
---: powdery mildew, fungicidal control of Erysiphe graminis in Indiana, 358
---: rust, natural rust epidemics in nurseries as affected by inoculum density, 410
---: ---: leaf: rust severity and host pathogen reaction types for wheat leaf rust in North Dakota in 1971, 77; severe leaf rust on wheat in Georgia, 534; 'Transfer', formerly "universally resistant" to all known natural uredial populations of Puccinia recondita in North America is susceptible to an Indiana isolate, 827; an inoculating system to inoculate seedlings with four uredial cultures of leaf rust simultaneously, 847
---: ---: stem: races of Puccinia graminis f. sp. tritici in the USA in 1971, 554; stem rust races in the Yaqui Valley of Mexico in 1972, 1038
---: ---: stripe, epiphytology of Puccinia striiformis in Oregon in 1971, 1071
---: Septoria leaf blotch: five spring wheat varieties tested for tolerance to Septoria nodorum, 188; 30 spring wheat varieties showed varied reactions to S. nodorum, 761; effect of Septoria leaf blotch on yield of spring wheat in Israel, 983
---: smut, chemical control of flag smut by seed treatment with oxycarboxin, carboxin, 2-methyl-5,6-dihydro-4H-pyrane-3-carbonic acid anilide and benomyl, 636
Wilt: of poinsettia, caused by lime in pot mix for potted plants, 415; Goss's wilt and leaf blight of corn in Nebraska, caused by a Corynebacterium, 863
---: bacterial wilt of cucumber, Erwinia tracheiphila, movement in susceptible and resistant cucumbers, 949
---: Fusarium: cyclamen wilt, caused by Fusarium oxysporum f. cyclaminis, in California, 493; of 'Dwarf Cavendish' banana in the Canary Islands, 1000; of watermelon, excellent wilt resistance of 'Calhoun Gray' and 'Summit' should be valuable to plant breeders, 1022
---: ---: of cotton: cotton loss by State in 1971, 328; response of 16 varieties of Egyptian cotton to wilt studied in soil infested with fungus alone or in combination with reniform nematode, 1040
---: ---: of tomato: an increase in soil pH from 6.0 to 7.0-7.5 decreased the incidence of Fusarium wilt, 151; development of resistant tomato varieties for control, 695; fumigation with DD-MENCS was greatly enhanced by paper or plastic mulches for control of Fusarium wilt, 953
---: persimmon wilt, caused by Cephalosporium diospyri, in Mississippi, 916
---: Verticillium: of tomato, increase in soil pH from 6.0 to 7.0-7.5 also increased the incidence of Verticillium wilt, 151; Verticillium appears to be major wilt problem of vegetables in Greece, 264; estimated cotton loss from Verticillium in 1971, 328
Wisconsin: 99; 712; 744; 798; 875
Wounding: apparently predisposes Madrone trees to new canker infection, 50

Xanthomonas carotae: 744
--- dieffenbachiae: 394; 904
--- malvacearum: 328; 765
--- phaseoli: 378
--- --- var. fuscans: 378
--- vesicatoria: 504
Xanthosoma spp.: dasheen mosaic virus was not inactivated in Xanthosoma corms below 65°C and above that temperature the corms did not germinate, 320
Xiphinema americanum: 476
--- index: 1031

Yellow poplar (Liriodendron tulipifera): incidence of Fusarium canker in South Carolina, 310
Yellows: of dandelion, mycoplasmalike bodies in phloem cells of yellows-infected plants, 270
Yield: hay and seed yields of 'Ranger' alfalfa were increased and leaf diseases were controlled with fungicides, 286; effect of Septoria leaf blotch on yield of spring wheat in Israel, 983
Yield loss: of cotton from parasitic diseases in 1971, 328; of alfalfa, from Phytophthora root rot in Arizona, 472; of soybeans inoculated with brown stem rot pathogen, Cephalosporium gregatum, 580; in early crop of Spanish peanuts in south Texas in 1971, 873; crown rot, leaf and scape disease of statice is a limiting factor in the production of yellow statice in Florida and causes losses in pink, white and blue varieties, 894

Zambia: 820; 1047; 1049
Zea mays (see Corn)
Zebra disease: resistance in Agave, and variation in Phytophthora isolates, 678

Prepared by Marguerite G. Benedict

U.S.Government Printing Office: 1973--511-883/13 Region No. 4

TRADE NAMES ARE USED IN THIS PUBLICATION SOLELY TO PROVIDE SPECIFIC INFORMATION. MENTION OF A TRADE NAME DOES NOT CONSTITUTE A WARRANTY OF THE PRODUCT BY THE U. S. DEPARTMENT OF AGRICULTURE OR AN ENDORSEMENT BY THE DEPARTMENT OVER OTHER PRODUCTS NOT MENTIONED.

IF PESTICIDES ARE HANDLED, APPLIED, OR DISPOSED OF IMPROPERLY, THEY MAY BE INJURIOUS TO HUMANS, DOMESTIC ANIMALS, DESIRABLE PLANTS, AND POLLINATING INSECTS, FISH, OR OTHER WILDLIFE, AND MAY CONTAMINATE WATER SUPPLIES. USE PESTICIDES ONLY WHEN NEEDED AND HANDLE THEM WITH CARE. FOLLOW THE DIRECTIONS AND HEED ALL PRECAUTIONS ON THE CONTAINER LABEL.

UNITED STATES DEPARTMENT OF AGRICULTURE
Agricultural Research Service
Beltsville, Maryland 20705

Official Business
Penalty for Private Use, $300

POSTAGE & FEES PAID
United States Department of Agriculture

PDR UNIVM300UG R 1
UNIV MICROFILMS
J REED SUPVR PERIODCALS
300 N ZEEB RD
ANN ARBOR MI 48106